

DYNC1LI2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP5478c

Product Information

Application IHC-P, WB, E **Primary Accession** 043237

Other Accession Q4R5P6, NP_006132.1

Reactivity Human **Predicted** Monkey Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB26902 54099 **Calculated MW** 152-180 **Antigen Region**

Additional Information

Gene ID 1783

Other Names Cytoplasmic dynein 1 light intermediate chain 2, Dynein light intermediate

chain 2, cytosolic, LIC-2, LIC53/55, DYNC1LI2, DNCLI2, LIC2

Target/SpecificityThis DYNC1LI2 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 152-180 amino acids from the Central

region of human DYNC1LI2.

Dilution IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions DYNC1LI2 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name DYNC1LI2 (HGNC:2966)

Synonyms DNCLI2, LIC2

Function Acts as one of several non-catalytic accessory components of the

cytoplasmic dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. May play a role in binding

dynein to membranous organelles or chromosomes.

Cellular Location Cytoplasm, cytoskeleton


Background

Cytoplasmic dynein is a microtubule-associated motor protein (Hughes et al., 1995 [PubMed 7738094]). See DYNC1H1 (MIM 600112) for general information about dyneins.


References

Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007) Olsen, J.V., et al. Cell 127(3):635-648(2006) Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)

Images



All lanes: Anti-DYNC1LI2 Antibody (Center) at 1:2000 dilution Lane 1: Human brain lysate Lane 2: U-87 MG whole cell lysate Lane 3: SK-OV-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 54 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



DYNC1LI2 Antibody (Center) (Cat. #AP5478c) immunohistochemistry analysis in formalin fixed and paraffin embedded human cervix carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the DYNC1LI2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.