10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# PREI3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54807

## Product Information

| | |
|---|---|
| **Application** | WB, IHC-P, IHC-F, IF, ICC, E |
| **Primary Accession** | Q9Y3A3 |
| **Reactivity** | Rat, Pig, Bovine |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Calculated MW** | 26032 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 25843 |
| **Other Names** | MOB-like protein phocein, 2C4D, Class II mMOB1, Mob1 homolog 3, Mob3, Mps one binder kinase activator-like 3, Preimplantation protein 3, MOB4, MOB3, MOBKL3, PHOCN, PREI3 |
| **Dilution** | WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000 |
| **Storage** | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

## Protein Information

| | |
|---|---|
| **Name** | MOB4 ( HGNC:17261) |
| **Function** | Part of the striatin-interacting phosphatase and kinase (STRIPAK) complexes. STRIPAK complexes have critical roles in protein (de)phosphorylation and are regulators of multiple signaling pathways including Hippo, MAPK, nuclear receptor and cytoskeleton remodeling. Different types of STRIPAK complexes are involved in a variety of biological processes such as cell growth, differentiation, apoptosis, metabolism and immune regulation. |
| **Cellular Location** | Cytoplasm, perinuclear region. Membrane; Peripheral membrane protein. Golgi apparatus, Golgi stack membrane; Peripheral membrane protein. Note=In a perinuclear punctate pattern. Associated with membranes and the Golgi stacks |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.