

VPREB1 Rabbit pAb

VPREB1 Rabbit pAb

Catalog # AP54826

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	P12018
Predicted	Human, Mouse, Rat, Pig, Horse, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16605
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from Human VPREB1/CD179a
Epitope Specificity	61-145/145
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the immunoglobulin superfamily. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
SUBUNIT	Associates non-covalently with IGLL1.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	VpreB is expressed in pre-B lymphocytes, but not in mature B cells or in other blood cell lineages. The gene which encodes VpreB maps to human chromosome 22q11.2. The VpreB and Lambda 5 genes encode the Iota and Omega polypeptide chains, respectively, which associate with the Ig-Mu chain to form a molecular complex that is expressed on the surface of pre-B cells. This complex presumably regulates Ig gene rearrangements in the early steps of B cell differentiation. In the mouse the two genes are simultaneously expressed in pre-B cells and belong to the same transcription unit. A primary transcript is synthesized from which the pre-B and Lambda 5 mRNAs are derived by alternative splicing. In the human, however, the two genes are separate and do not belong to the same transcription unit.

Additional Information

Gene ID	7441
Other Names	Immunoglobulin iota chain, CD179 antigen-like family member A, Protein VPreB1, V(pre)B protein, VpreB protein, CD179a, VPREB1, VPREB
Target/Specificity	Only expressed by pre-B-cells.
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	VPREB1
Synonyms	VPREB
Function	Associates with the Ig-mu chain to form a molecular complex that is expressed on the surface of pre-B-cells. This complex presumably regulates Ig gene rearrangements in the early steps of B- cell differentiation.
Cellular Location	Endoplasmic reticulum {ECO:0000250 UniProtKB:P13372}
Tissue Location	Only expressed by pre-B-cells.

Background

VpreB is expressed in pre-B lymphocytes, but not in mature B cells or in other blood cell lineages. The gene which encodes VpreB maps to human chromosome 22q11.2. The VpreB and Lambda 5 genes encode the Iota and Omega polypeptide chains, respectively, which associate with the Ig-Mu chain to form a molecular complex that is expressed on the surface of pre-B cells. This complex presumably regulates Ig gene rearrangements in the early steps of B cell differentiation. In the mouse the two genes are simultaneously expressed in pre-B cells and belong to the same transcription unit. A primary transcript is synthesized from which the pre-B and Lambda 5 mRNAs are derived by alternative splicing. In the human, however, the two genes are separate and do not belong to the same transcription unit.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.