

ARIH2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54877

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O95376
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57819
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ARIH2
Epitope Specificity	41-140/493
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Cytoplasm.
SIMILARITY	Belongs to the RBR family. Ariadne subfamily. Contains 1 IBR-type zinc finger. Contains 2 RING-type zinc fingers.
SUBUNIT	Interacts (via RING-type 1) with UBE2L3. Interacts (via RING-type 2) with UBE2N. Interacts (via RING-type 2) with GFI1B. Interacts with GFI1; prevents its ubiquitination and proteasomal degradation.
Post-translational modifications	Ubiquitinated. Ubiquitination promotes proteasomal degradation. Phosphorylated upon DNA damage, probably by ATM or ATR.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	TRIAD1, also known as ARIH2 (ariadne homolog 2) or ARI2, is a 493 amino acid protein that contains one IBR-type zinc finger and two RING-type zinc fingers and belongs to the ariadne subfamily of RBR proteins. Localized to the nucleus, TRIAD1 interacts with UBE2L3 and is thought to act as an E3 ubiquitin-protein ligase, functioning to accept ubiquitin from E2 ubiquitin-conjugating enzymes and transfer the acquired ubiquitin residue to target substrates. TRIAD1 is subject to post-translational DNA damage-dependent phosphorylation, probably by ATM or ATR. The gene encoding TRIAD1 maps to human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci.

Additional Information


Gene ID	10425
Other Names	E3 ubiquitin-protein ligase ARIH2, ARI-2, Protein ariadne-2 homolog, 2.3.2.31, RING-type E3 ubiquitin transferase ARIH2, Triad1 protein, ARIH2, ARI2, TRIAD1 {ECO:0000303 PubMed:16118314}

Target/Specificity	Widely expressed with higher expression in granulocytes.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ARIH2 {ECO:0000303 PubMed:31253590, ECO:0000312 HGNC:HGNC:690}
Function	E3 ubiquitin-protein ligase, which catalyzes ubiquitination of target proteins together with ubiquitin-conjugating enzyme E2 UBE2L3 (PubMed: 16118314 , PubMed: 17646546 , PubMed: 19340006 , PubMed: 24076655 , PubMed: 33268465 , PubMed: 34518685 , PubMed: 38418882). Acts as an atypical E3 ubiquitin-protein ligase by working together with cullin-5-RING ubiquitin ligase complex (ECS complex, also named CRL5 complex) and initiating ubiquitination of ECS substrates: associates with ECS complex and specifically mediates addition of the first ubiquitin on ECS targets (PubMed: 33268465 , PubMed: 34518685 , PubMed: 38418882). The initial ubiquitin is then elongated (PubMed: 33268465). E3 ubiquitin- protein ligase activity is activated upon binding to neddylated form of the cullin-5 (CUL5) component of the ECS complex (PubMed: 24076655). Together with the ECS(ASB9) complex, catalyzes ubiquitination of CKB (PubMed: 33268465). Promotes ubiquitination of DCUN1D1 (PubMed: 30587576). Mediates 'Lys-6', 'Lys-48'- and 'Lys-63'-linked polyubiquitination (PubMed: 16118314 , PubMed: 17646546 , PubMed: 19340006). May play a role in myelopoiesis (PubMed: 19340006).
Cellular Location	Nucleus. Cytoplasm
Tissue Location	Widely expressed with higher expression in granulocytes.

Images



Sample:
Pancreas (Mouse) Lysate at 40 ug
Primary: Anti-ARIH2 (AP54877) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 58 kD
Observed band size: 58 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.