10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ARVCF Rabbit pAb

ARVCF Rabbit pAb
Catalog # AP54882

Product Information

Application	WB
Primary Accession	O00192
Reactivity	Rat, Mouse
Predicted	Pig, Dog, Human, Rabbit, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	104642
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ARVCF
Epitope Specificity	501-600/962
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Armadillo Repeat gene deleted in Velo-Cardio-Facial syndrome (ARVCF) is a member of the catenin family. This family plays an important role in the formation of adherens junction complexes, which are thought to facilitate communication between the inside and outside environments of a cell. The ARVCF gene was isolated in the search for the genetic defect responsible for the autosomal dominant Velo-Cardio-Facial syndrome (VCFS), a relatively common human disorder with phenotypic features including cleft palate, conotruncal heart defects and facial dysmorphology. The ARVCF gene encodes a protein containing two motifs, a coiled coil domain in the N-terminus and a 10 armadillo repeat sequence in the midregion. Since these sequences can facilitate protein-protein interactions ARVCF is thought to function in a protein complex. In addition, ARVCF contains a predicted nuclear-targeting sequence suggesting that it may have a function as a nuclear protein. [provided by RefSeq, Jun 2010].

Additional Information

Gene ID	421
Other Names	Splicing regulator ARVCF, Armadillo repeat protein deleted in velo-cardio-facial syndrome, ARVCF (HGNC:728)
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name	ARVCF (HGNC:728)
Function	Contributes to the regulation of alternative splicing of pre- mRNAs.
Cellular Location	Cell junction, adherens junction. Nucleus. Cytoplasm Note=In heart, localizes at area composita, the mixed-type junctional structure composed of both desmosomal and adherens junctional proteins {ECO:0000250|UniProtKB:B4F7F3}
Tissue Location	Found in all the examined tissues including heart, brain, liver and kidney. Found at low level in lung. Expressed in dermal connective tissue, salivary gland duct and in the corneal layer (at protein level) (PubMed:30479852). Expressed in arrector pili muscle (at protein level) (PubMed:29034528). High levels detected in epithelial cells with lower levels found in fibroblasts and T lymphocytes (PubMed:10725230).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:
Cerebrum (Mouse) Lysate at 40 ug
Cerebrum (Rat) Lysate at 40 ug
Primary: Anti- ARVCF (AP54882) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 105 kD
Observed band size: 105 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.