

Attractin Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54885

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	O75882
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	158537

Additional Information

Gene ID	8455
Other Names	Attractin, DPPT-L, Mahogany homolog, ATRN, KIAA0548, MGCA
Dilution	WB=1:500-2000, Elisa=1:500-1000, IHC-P=1:100-500, IHC-F=1:100-500, IF=1:100-500, ICC=1:100-500,
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ATRN
Synonyms	KIAA0548, MGCA
Function	Involved in the initial immune cell clustering during inflammatory response and may regulate chemotactic activity of chemokines. May play a role in melanocortin signaling pathways that regulate energy homeostasis and hair color. Low-affinity receptor for agouti (By similarity). Has a critical role in normal myelination in the central nervous system (By similarity).
Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 3]: Secreted
Tissue Location	Isoform 2 is detected in plasma (at protein level). Expressed and secreted by activated T-lymphocytes. Expressed at low to moderate levels in peripheral blood leukocytes, spleen, lymph node, tonsil, bone marrow and fetal liver. At very low levels found in thymus. Isoform 2 is the major isoform in peripheral blood leukocytes

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.