

BDP Rabbit pAb

BDP Rabbit pAb
Catalog # AP54892

Product Information

Application	WB
Reactivity	Human
Predicted	Mouse, Rat, Dog, Horse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human BDP/ARID3B
Epitope Specificity	251-350/561
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
SIMILARITY	Contains 1 ARID domain. Contains 1 REKLES domain.
SUBUNIT	Heterodimer with ARID3A. Interacts with unphosphorylated RB1.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	ARID3B represents a member of a unique family of DNA-binding proteins that have roles in cell lineage gene regulation, cell cycle control, embryonic patterning and transcriptional regulation. ARID3B localizes to the nucleus and contains an AT-rich interactive domain (ARID) which has DNA-binding capabilities. Functioning as a transcription factor, ARID3B associates in a heterodimer with the related protein ARID3A and is believed to play an important role in neural crest survival during embryogenesis. In addition, ARID3B may participate in malignant transformation and neuroblastoma growth, suggesting a possible use of ARID3B as a tumor marker for neuroblastoma.

Additional Information


Other Names	AT-rich interactive domain-containing protein 3B, ARID domain-containing protein 3B, Bright and dead ringer protein, Bright-like protein, ARID3B, BDP, DRIL2
Target/Specificity	Expressed in placenta, testis and leukocytes. Expressed in neuroblastoma. Present in K562 erythrocytic leukemia cells (at protein level).
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Background

ARID3B represents a member of a unique family of DNA-binding proteins that have roles in cell lineage gene regulation, cell cycle control, embryonic patterning and transcriptional regulation. ARID3B localizes to the nucleus and contains an AT-rich interactive domain (ARID) which has DNA-binding capabilities. Functioning as a transcription factor, ARID3B associates in a heterodimer with the related protein ARID3A and is believed to play an important role in neural crest survival during embryogenesis. In addition, ARID3B may participate in malignant transformation and neuroblastoma growth, suggesting a possible use of ARID3B as a tumor marker for neuroblastoma.

Images



Sample:

Lane 1: K562 (Human) Cell Lysate at 30 ug

Lane 2: SH-SY5Y (Human) Cell Lysate at 30 ug

Lane 3: HepG2 (Human) Cell Lysate at 30 ug

Lane 4: Molt-4 (Human) Cell Lysate at 30 ug

Primary:

Anti-BDP (AP54892) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 61 kD

Observed band size: 61 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.