

PEX19 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP54906

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	P40855
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32807

Additional Information

Gene ID	5824
Other Names	Peroxisomal biogenesis factor 19, 33 kDa housekeeping protein, Peroxin-19, Peroxisomal farnesylated protein, PEX19, HK33, PXF
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	PEX19 (HGNC:9713)
Synonyms	HK33, PXF
Function	Necessary for early peroxisomal biogenesis. Acts both as a cytosolic chaperone and as an import receptor for peroxisomal membrane proteins (PMPs). Binds and stabilizes newly synthesized PMPs in the cytoplasm by interacting with their hydrophobic membrane-spanning domains, and targets them to the peroxisome membrane by binding to the integral membrane protein PEX3. Excludes CDKN2A from the nucleus and prevents its interaction with MDM2, which results in active degradation of TP53.
Cellular Location	Cytoplasm. Peroxisome membrane; Lipid-anchor; Cytoplasmic side. Note=Mainly cytoplasmic. Some fraction membrane-associated to the outer surface of peroxisomes.
Tissue Location	Ubiquitously expressed. Isoform 1 is strongly predominant in all tissues

except in utero where isoform 2 is the main form.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.