

ZNF207 Rabbit pAb

ZNF207 Rabbit pAb

Catalog # AP54975

Product Information

Application	WB
Primary Accession	O43670
Reactivity	Human, Mouse
Predicted	Rat, Pig, Dog, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50751
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ZNF207
Epitope Specificity	321-420/478
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	7756
Other Names	BUB3-interacting and GLEBS motif-containing protein ZNF207, BuGZ, hBuGZ, Zinc finger protein 207, ZNF207 (HGNC:12998)
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ZNF207 (HGNC:12998)
Function	Kinetochores- and microtubule-binding protein that plays a key role in spindle assembly (PubMed: 24462186 , PubMed: 24462187 , PubMed: 26388440). ZNF207/BuGZ is mainly composed of disordered low-complexity regions and undergoes phase transition or coacervation to form temperature-dependent liquid droplets. Coacervation promotes microtubule bundling and concentrates tubulin, promoting microtubule polymerization

and assembly of spindle and spindle matrix by concentrating its building blocks (PubMed:[26388440](#)). Also acts as a regulator of mitotic chromosome alignment by mediating the stability and kinetochore loading of BUB3 (PubMed:[24462186](#), PubMed:[24462187](#)). Mechanisms by which BUB3 is protected are unclear: according to a first report, ZNF207/BuGZ may act by blocking ubiquitination and proteasomal degradation of BUB3 (PubMed:[24462186](#)). According to another report, the stabilization is independent of the proteasome (PubMed:[24462187](#)).

Cellular Location

Nucleus. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, spindle. Note=Localizes primarily to the nucleus in interphase, concentrates at kinetochores prior to nuclear envelope breakdown and during early prometaphase, and disappears from kinetochores upon microtubule-binding


Tissue Location

Ubiquitous..

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:

Embryo (Mouse) Lysate at 40 ug

HeLa (Human) Lysate at 30 ug

Primary: Anti-ZNF207 (AP54975) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 51 kD

Observed band size: 51 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.