

DMAP1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55040

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9NPF5
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52993
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DMAP1
Epitope Specificity	31-130/467
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Cytoplasm. Note=Targeted to replication foci throughout S phase by DNMT1.
SIMILARITY	Contains 1 SANT domain.
SUBUNIT	Component of the NuA4 histone acetyltransferase complex which contains the catalytic subunit KAT5/TIP60 and the subunits EP400, TRRAP/PAF400, BRD8/SMAP, EPC1, DMAP1/DNMAP1, RUVBL1/TIP49, RUVBL2, ING3, actin, ACTL6A/BAF53A, MORF4L1/MRG15, MORF4L2/MRGX, MRGBP, YEATS4/GAS41, VPS72/YL1 and MEAF6. Component of a NuA4-related complex which contains EP400, TRRAP/PAF400, SRCAP, BRD8/SMAP, EPC1, DMAP1/DNMAP1, RUVBL1/TIP49, RUVBL2, actin, ACTL6A/BAF53A, VPS72 and YEATS4/GAS41. DMAP1 also forms a complex with DNMT1 and HDAC2. Throughout S phase it interacts directly with the N-terminus of DNMT1, which serves to recruit DMAP1 to replication foci. DMAP1 interacts with ING1, a component of the mSin3A transcription repressor complex, although this interaction is not required for recruitment of ING1 to heterochromatin. Interacts directly with the transcriptional corepressor TSG101. Interacts with the pro-apoptotic protein DAXX. Interacts with URI1.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Methylation of DNA contributes to the regulation of gene transcription in eukaryotic systems. DNA methylation is predominantly found on cytosine residues that are present in dinucleotide motifs consisting of a 5' cytosine followed by a guanosine (CpG), and it requires the enzymatic activity of DNA methyltransferases (DNMTs), which results in transcriptional repression of the methylated gene. DNA methyltransferase 1-associating protein (Dmap1) binds to methyl-CpG rich domains and mediate the transcriptional inhibition associated with DNA methylation. Dmap1 interacts with Daxx to enhanced Daxx-mediated repression of glucocorticoid receptor transcriptional activity. Daxx also protects Dmap1 from protein degradation in vivo.

Additional Information

Gene ID	55929
Other Names	DNA methyltransferase 1-associated protein 1, DNMAP1, DNMT1-associated protein 1, DMAP1, KIAA1425
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DMAP1
Synonyms	KIAA1425
Function	Involved in transcription repression and activation. Its interaction with HDAC2 may provide a mechanism for histone deacetylation in heterochromatin following replication of DNA at late firing origins. Can also repress transcription independently of histone deacetylase activity. May specifically potentiate DAXX-mediated repression of glucocorticoid receptor-dependent transcription. Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage. Participates in the nuclear localization of URI1 and increases its transcriptional corepressor activity.
Cellular Location	Nucleus. Cytoplasm. Note=Targeted to replication foci throughout S phase by DNMT1

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.