10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

DNAJC9 Rabbit pAb

DNAJC9 Rabbit pAb
Catalog # AP55041

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q8WXX5
Reactivity	Rat, Mouse
Predicted	Pig, Dog, Human, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29910
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DNAJC9
Epitope Specificity	161-260/260
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The DnaJ family comprises a group of chaperone proteins that contain a J domain and have diverse cellular localization and functions. DnaJ proteins play a critical role in the HSP 70 chaperone machine by interacting with HSP 70 to stimulate ATP hydrolysis and are also important mediators of proteolysis and protein degradation. DnaJC9 (DnaJ (Hsp40) homolog, subfamily C, member 9), also designated HDJC9, JDD1 or DnaJ protein SB73, is a 260 amino acid protein found at moderate levels in most tissues with highest expression in the germinal zone of the central nervous system, testis, ovary, renal cortex and fetal liver. A member of the DnaJ family, DnaJC9 contains one N-terminal J domain but lacks the typical G/F and zinc finger regions that are typical of DnaJ family members. DnaJC9 localiizes to nuclei under normal conditions but may be transported to cytoplasm and plasma membrane when exposed to heat shock.

Additional Information

Gene ID	23234
Other Names	DnaJ homolog subfamily C member 9, HDJC9, DnaJ protein SB73, DNAJC9
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	DNAJC9
Function	Acts as a dual histone chaperone and heat shock co-chaperone (PubMed:33857403). As a histone chaperone, forms a co-chaperone complex with MCM2 and histone H3-H4 heterodimers; and may thereby assist MCM2 in histone H3-H4 heterodimer recognition and facilitate the assembly of histones into nucleosomes (PubMed:33857403). May also act as a histone co-chaperone together with TONSL (PubMed:33857403). May recruit histone chaperones ASF1A, NASP and SPT2 to histone H3-H4 heterodimers (PubMed:33857403). Also plays a role as co-chaperone of the HSP70 family of molecular chaperone proteins, such as HSPA1A, HSPA1B and HSPA8 (PubMed:17182002, PubMed:33857403). As a co-chaperone, may play a role in the recruitment of HSP70-type molecular chaperone machinery to histone H3-H4 substrates, thereby maintaining the histone structural integrity (PubMed:33857403). Exhibits activity to assemble histones onto DNA in vitro (PubMed:33857403).
Cellular Location	Nucleus. Cytoplasm. Cell membrane. Note=Predominantly nuclear. Translocates to the cytoplasm and membrane after heat shock
Tissue Location	Expressed in heart, placenta, liver, skeletal muscle, kidney, pancreas, thymus, ovary, colon and peripheral blood


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:
Thymus (Mouse) Lysate at 40 ug
Primary: Anti- DNAJC9 (AP55041) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 30 kD
Observed band size: 35 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DNAJC9) Polyclonal Antibody, Unconjugated (AP55041) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.