

DPP3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55042

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9NY33
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	82589
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DPP3
Epitope Specificity	1-100/737
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm.
SIMILARITY	Belongs to the peptidase M49 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Dipeptidyl peptidases (DPPs) mediate regulatory activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs can bind specific voltage-gated potassium channels and alter their expression and biophysical properties and may also influence T cells. DPP proteins include DPRP1, DPRP2, DPP3, DPP7, DPP10, DPPX and CD26. DPP3 (dipeptidyl-peptidase 3), also known as DPPIII, is a zinc-exopeptidase that belongs to the peptidase M49 family. DPP3 localizes to the cytoplasm and is involved in intracellular protein catabolism. More specifically, DPP3 is an important enzyme involved in the degradation of enkephalins. An increase in the activity of DPP3 is implicated in ovarian and endometrial cancers.

Additional Information

Gene ID	10072
Other Names	Dipeptidyl peptidase 3, 3.4.14.4, Dipeptidyl aminopeptidase III, Dipeptidyl arylamidase III, Dipeptidyl peptidase III, DPP III, Enkephalinase B, DPP3
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	DPP3
Function	Cleaves and degrades bioactive peptides, including angiotensin, Leu-enkephalin and Met-enkephalin (PubMed: 1515063 , PubMed: 3233187). Also cleaves Arg-Arg-beta-naphthylamide (in vitro) (PubMed: 11209758 , PubMed: 3233187 , PubMed: 9425109).
Cellular Location	Cytoplasm, cytosol
Tissue Location	Detected in placenta (at protein level) (PubMed:3233187). Detected in erythrocytes (at protein level) (PubMed:1515063).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.