

DYX1C1 Rabbit pAb

DYX1C1 Rabbit pAb

Catalog # AP55046

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q8WXU2
Reactivity	Rat, Mouse
Predicted	Dog, Human, Zebrafish, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48527
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DYX1C1
Epitope Specificity	51-120/420
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Cytoplasm.
DISEASE	Defects in DYX1C1 may be a cause of susceptibility to dyslexia type 1 (DYX1) [MIM:127700]. A relatively common, complex cognitive disorder characterized by an impairment of reading performance despite adequate motivational, educational and intellectual opportunities. It is a multifactorial trait, with evidence for familial clustering and heritability. Note=A chromosomal aberration involving DYX1C1 has been found in a family affected by dyslexia. Translocation t(2;15)(q11;q21).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The gene encoding DYX1C1 maps in the 15q21 region, which is disrupted by a translocation t(2;15)(q11;q21) and segregates with dyslexia. Two sequence changes in DYX1C1, including one involving the translation initiation sequence and an Elk-1 transcription factor binding site (-3G --> A) and a codon (1249G --> T), introduce a premature stop codon and truncate the protein by 4 amino acids. DYX1C1 encodes a nuclear tetratricopeptide repeat domain protein that is dynamically regulated in brain. In human brain, DYX1C1 protein localizes to a fraction of cortical neurons and white matter glial cells. It is also expressed in lung, kidney and testis.

Additional Information

Gene ID	161582
Other Names	Dynein axonemal assembly factor 4 {ECO:0000312 HGNC:HGNC:21493}, Dyslexia susceptibility 1 candidate gene 1 protein, DNAAF4 (HGNC:21493), DYX1C1, EKN1

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	DNAAF4 (HGNC:21493)
Synonyms	DYX1C1, EKN1
Function	Axonemal dynein assembly factor required for ciliary motility. Involved in neuronal migration during development of the cerebral neocortex. May regulate the stability and proteasomal degradation of the estrogen receptors that play an important role in neuronal differentiation, survival and plasticity.
Cellular Location	Nucleus. Cytoplasm. Dynein axonemal particle. Cell projection, neuron projection {ECO:0000250 UniProtKB:Q5VJS5}
Tissue Location	Expressed in several tissues, including brain, lung, kidney and testis. In brain localizes to a fraction of cortical neurons and white matter glial cells.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:
Kidney (Mouse) Lysate at 40 ug
Primary: Anti- DYX1C1 (AP55046) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 49 kD
Observed band size: 49 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DYX1C1) Polyclonal Antibody, Unconjugated (AP55046) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.