

ECHDC2 Rabbit pAb

ECHDC2 Rabbit pAb
Catalog # AP55049

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q86YB7
Predicted	Human, Mouse, Rat, Pig, Horse, Zebrafish, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31126
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ECHDC2
Epitope Specificity	153-260/292
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Mitochondrion (Potential).
SIMILARITY	Belongs to the enoyl-CoA hydratase/isomerase family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	ECHDC2 is a 292 amino acid mitochondrial protein that exists as two isoforms which are produced by alternative splicing events. The gene encoding ECHDC2 maps to chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 encodes a large number of disease-associated proteins, including Lamin A which, when expressed abnormally, can build up in the nucleus and cause nuclear blebs, a characteristic of the rare aging disease Hutchinson-Gilford progeria. Additionally, genes that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinsons Disease, Gaucher disease, schizophrenia and Usher syndrome are all located on chromosome 1. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

Additional Information

Gene ID	55268
Other Names	Enoyl-CoA hydratase domain-containing protein 2, mitochondrial, ECHDC2
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ECHDC2
Cellular Location	Mitochondrion.

Background

ECHDC2 is a 292 amino acid mitochondrial protein that exists as two isoforms which are produced by alternative splicing events. The gene encoding ECHDC2 maps to chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 encodes a large number of disease-associated proteins, including Lamin A which, when expressed abnormally, can build up in the nucleus and cause nuclear blebs, a characteristic of the rare aging disease Hutchinson-Gilford progeria. Additionally, genes that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinsons Disease, Gaucher disease, schizophrenia and Usher syndrome are all located on chromosome 1. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.