

GBP1 Rabbit pAb

GBP1 Rabbit pAb
Catalog # AP55125

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P32455
Predicted	Rat, Pig, Dog, Human, Mouse, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67931
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GBP1
Epitope Specificity	21-120/592
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Golgi apparatus membrane; Lipid-anchor; Cytoplasmic side. Secreted. Note=Secreted from endothelial cells in the cerebrospinal fluid, upon bacterial challenge and independently of interferon-gamma induction. Golgi membrane localization requires isoprenylation and the presence of another IFN-gamma-induced factor.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Guanylate-binding proteins, GBP1 and GBP2 are GTP-binding proteins with a high-turnover GTPase activity and an antiviral effect (1-4). GBP1 mediates an antiviral effect against vesicular stomatitis virus and encephalomyocarditis virus and plays a role in the IFN-mediated antiviral response against these viruses (4). GBP1 and GBP2 belong to a group of large GTP-binding proteins with a high concentration-dependent GTPase activity that have the common ability to undergo oligomerization (1). GBP1 and GBP2 are bone marrow-derived GTPases encoded by interferon-activated genes and are inducible following IFN treatment (2,3). Specifically, GBP1 is expressed in cultured mammary epithelial tumor cell lines after treatment with IFN-gamma and LPS.

Additional Information

Gene ID	2633
Other Names	Guanylate-binding protein 1 {ECO:0000303 PubMed:39394410, ECO:0000312 HGNC:HGNC:4182}, 3.6.1.-, 3.6.5.-, GTP-binding protein 1, GBP-1, HuGBP-1, hGBP1, Guanine nucleotide-binding protein 1, Interferon-induced guanylate-binding protein 1, GBP1 {ECO:0000303 PubMed:7512561, ECO:0000312 HGNC:HGNC:4182}

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	GBP1 {ECO:0000303 PubMed:7512561, ECO:0000312 HGNC:HGNC:4182}
Function	Interferon (IFN)-inducible GTPase that plays important roles in innate immunity against a diverse range of bacterial, viral and protozoan pathogens (PubMed:16511497, PubMed:22106366, PubMed:29144452, PubMed:31268602, PubMed:32510692, PubMed:32581219, PubMed:37797010, PubMed:7512561). Hydrolyzes GTP to GMP in two consecutive cleavage reactions: GTP is first hydrolyzed to GDP and then to GMP in a processive manner (PubMed:16511497, PubMed:32510692, PubMed:7512561, PubMed:39394410). Following infection, recruited to the pathogen- containing vacuoles or vacuole-escaped bacteria and promotes both inflammasome assembly and autophagy (PubMed:29144452, PubMed:31268602). Acts as a positive regulator of inflammasome assembly by facilitating the detection of inflammasome ligands from pathogens (PubMed:31268602, PubMed:32510692, PubMed:32581219). Involved in the lysis of pathogen- containing vacuoles, releasing pathogens into the cytosol (By similarity). Following pathogen release in the cytosol, forms a protein coat in a GTPase-dependent manner that encapsulates pathogens and promotes the detection of ligands by pattern recognition receptors (PubMed:32510692, PubMed:32581219). Plays a key role in inflammasome assembly in response to infection by Gram-negative bacteria: following pathogen release in the cytosol, forms a protein coat that encapsulates Gram-negative bacteria and directly binds to lipopolysaccharide (LPS), disrupting the O-antigen barrier and unmasking lipid A that is that detected by the non-canonical inflammasome effector CASP4/CASP11 (PubMed:32510692, PubMed:32581219). Also promotes recruitment of proteins that mediate bacterial cytolysis, leading to release double- stranded DNA (dsDNA) that activates the AIM2 inflammasome (PubMed:31268602). Involved in autophagy by regulating bacteriolytic peptide generation via its interaction with ubiquitin-binding protein SQSTM1, which delivers monoubiquitinated proteins to autolysosomes for the generation of bacteriolytic peptides (By similarity). Confers protection to several pathogens, including the bacterial pathogens <i>L.monocytogenes</i> and <i>M.bovis</i> BCG as well as the protozoan pathogen <i>T.gondii</i> (PubMed:31268602). Exhibits antiviral activity against influenza virus (PubMed:22106366).
Cellular Location	Cytoplasmic vesicle membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor; Cytoplasmic side. Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, cytosol. Secreted. Note=Localizes to pathogen-containing vacuoles or to the cell surface of bacteria that escaped vacuoles (PubMed:29144452, PubMed:31268602, PubMed:32510692, PubMed:32581219) Secreted from endothelial cells in the cerebrospinal fluid, upon bacterial challenge and independently of IFNG induction (PubMed:16936281). Golgi membrane localization requires isoprenylation and the presence of another IFNG-induced factor (PubMed:15937107) Sequestered in the cytosol following phosphorylation by PIM1 and subsequent interaction with 14-3-3 protein sigma (SFN) (PubMed:37797010).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.