

# Gemin 7 Rabbit pAb

Gemin 7 Rabbit pAb Catalog # AP55135

#### **Product Information**

**Application** IHC-P, IHC-F, IF

**Primary Accession Q9H840** 

Reactivity Human, Chicken, Dog

Host Rabbit Clonality Polyclonal 14537 Calculated MW **Physical State** Liquid

KLH conjugated synthetic peptide derived from human Gemin 7 **Immunogen** 

**Epitope Specificity** 21-120/131 Isotype IgG

**Purity** affinity purified by Protein A

**Buffer** 

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. SUBCELLULAR LOCATION Nucleus, nucleoplasm, Nucleus, gem, Cytoplasm, Note=Found both in the

nucleoplasm and in nuclear bodies called gems (Gemini of Cajal bodies) that are often in proximity to Cajal (coiled) bodies. Also found in the cytoplasm.

**SIMILARITY** Belongs to the gemin-7 family.

**SUBUNIT** Part of the core SMN complex that contains SMN1, GEMIN2/SIP1,

DDX20/GEMIN3, GEMIN4, GEMIN5, GEMIN6, GEMIN7, GEMIN8 and

STRAP/UNRIP. Interacts directly with GEMIN6, SNRPB, SNRPD2, SNRPD3 and

This product as supplied is intended for research use only, not for use in **Important Note** 

human, therapeutic or diagnostic applications.

**Background Descriptions** Gemin7 is a 131 amino acid protein encoded by the mouse gene Gemin7.

> Gemin7, along with Gemin6, is a significant component of the the large multiprotein human SMN complex. The SMN complex functions as an assembly machine for small nuclear ribonucleoproteins (snRNPs)-the major components of the spliceosome. The survival of motor neurons (SMN) protein, a product of the disease gene of the common neurodegenerative disease, spinal muscular atrophy, is also part of the SMN complex. Although Gemin6 and Gemin7 have no significant sequence similarity with Sm proteins, both adopt canonical Sm folds. Moreover, Gemin6 and Gemin7 exist as a heterodimer, and interact with each other via an interface similar to that which mediates interactions among the Sm proteins. The Gemin6/Gemin7 complex binds to Sm proteins and might help organize Sm proteins for

formation of Sm rings on snRNA targets.

### **Additional Information**

Gene ID 79760

**Other Names** Gem-associated protein 7, Gemin-7, SIP3, GEMIN7 **Dilution** 

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

#### **Protein Information**

**GEMIN7** Name

**Function** The SMN complex catalyzes the assembly of small nuclear

ribonucleoproteins (snRNPs), the building blocks of the spliceosome, and thereby plays an important role in the splicing of cellular pre- mRNAs. Most spliceosomal snRNPs contain a common set of Sm proteins SNRPB, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF and SNRPG that assemble in a heptameric protein ring on the Sm site of the small nuclear RNA to form the core snRNP (Sm core). In the cytosol, the Sm proteins SNRPD1, SNRPD2, SNRPE, SNRPF and SNRPG are trapped in an inactive 6S pICln-Sm complex by the chaperone CLNS1A that controls the assembly of the core snRNP. To assemble core snRNPs, the SMN complex accepts the trapped 5Sm proteins from CLNS1A forming an intermediate. Binding of snRNA inside 5Sm triggers eviction of the SMN complex, thereby allowing binding of SNRPD3 and SNRPB to complete

assembly of the core snRNP.


**Cellular Location** Nucleus, nucleoplasm. Nucleus, gem. Cytoplasm Note=Found both in the

nucleoplasm and in nuclear bodies called gems (Gemini of Cajal bodies) that are often in proximity to Cajal (coiled) bodies. Also found in the cytoplasm

**Background** 


Gemin7 is a 131 amino acid protein encoded by the mouse gene Gemin7. Gemin7, along with Gemin6, is a significant component of the the large multiprotein human SMN complex. The SMN complex functions as an assembly machine for small nuclear ribonucleoproteins (snRNPs)-the major components of the spliceosome. The survival of motor neurons (SMN) protein, a product of the disease gene of the common neurodegenerative disease, spinal muscular atrophy, is also part of the SMN complex. Although Gemin6 and Gemin7 have no significant sequence similarity with Sm proteins, both adopt canonical Sm folds. Moreover, Gemin6 and Gemin7 exist as a heterodimer, and interact with each other via an interface similar to that which mediates interactions among the Sm proteins. The Gemin6/Gemin7 complex binds to Sm proteins and might help organize Sm proteins for formation of Sm rings on snRNA targets.

## **Images**



Blank control: Hela(blue), the cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice.. Isotype Control Antibody: Rabbit IgG(orange); Secondary Antibody: Goat anti-rabbit IgG-FITC(white blue).


Dilution: 1:100 in 1 X PBS containing 0.5% BSA; Primary Antibody Dilution: 1 µg in 100 µL1X PBS containing 0.5% BSA(green).



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Gemin 7 Polyclonal Antibody,

Unconjugated(AP55135) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: Hela(blue), the cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice..

Isotype Control Antibody: Rabbit IgG(orange); Secondary Antibody: Goat anti-rabbit IgG-FITC(white blue), Dilution: 1:100 in 1 X PBS containing 0.5% BSA; Primary Antibody Dilution: 1 µg in 100 µL1X PBS containing 0.5% BSA(green).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.