

GPX2 Rabbit pAb

GPX2 Rabbit pAb
Catalog # AP55158

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P18283
Reactivity	Rat, Mouse
Predicted	Dog, Human, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21954
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GPX2/Glutathione Peroxidase 2
Epitope Specificity	41-140/190
Isotype	IgG
Purity	affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.**SUBCELLULAR LOCATION** Cytoplasm. Mainly cytoplasmic.**Important Note** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.**Background Descriptions** Glutathione peroxidase (GPx) enzymes are generally selenium-containing tetrameric glycoproteins that help prevent lipid peroxidation of cell membranes. GPx enzymes reduce lipid hydroperoxides to alcohols, and reduce free hydrogen peroxide to water. GPx members are among the few proteins known in higher vertebrates to contain selenocysteine, which occurs at the active site of glutathione peroxidase and is coded by the nonsense (stop) codon TGA. There are eight GPx homologs (GPx-1–8). GPx-1 plays an important role in the antioxidant defense of the vascular wall and neural cells in response to oxidative stress. GPx-2 is the major isoform in the lungs and its basal or inducible expression is dependent on Nrf2. GPx-3 is under regulation by hypoxic stress and the expression and deficiency of GPx-3 is associated with cardiovascular disease and stroke. GPx-5 is selenium-independent; it is bound to the acrosome of sperm, where it may protect sperm from premature acrosome reaction in the epididymis.

Additional Information

Gene ID 2877**Other Names** Glutathione peroxidase 2, GPx-2, GSHPx-2, 1.11.1.9, Gastrointestinal glutathione peroxidase, Glutathione peroxidase-gastrointestinal, GPx-GI, GSHPx-GI, Glutathione peroxidase-related protein 2, GPRP-2, Phospholipid hydroperoxide glutathione peroxidase GPX2, 1.11.1.12, GPX2 {ECO:0000303 | Ref.9, ECO:0000312 | HGNC:HGNC:4554}

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	GPX2 {ECO:0000303 Ref.9, ECO:0000312 HGNC:HGNC:4554}
Function	Catalyzes the reduction of hydroperoxides in a glutathione- dependent manner thus regulating cellular redox homeostasis (PubMed: 36608588 , PubMed: 8428933). Can reduce small soluble hydroperoxides such as H ₂ O ₂ , cumene hydroperoxide and tert-butyl hydroperoxide, as well as several fatty acid-derived hydroperoxides (PubMed: 36608588 , PubMed: 8428933). Cannot reduce phosphatidylcholine hydroperoxide (PubMed: 8428933).
Cellular Location	Cytoplasm, cytosol.
Tissue Location	Mostly in liver and gastrointestinal tract, not found in heart or kidney.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:

Lane 1: Mouse Liver tissue lysates


Lane 2: Rat Kidney tissue lysates

Primary: Anti-GPX2 (AP55158) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 22 kDa

Observed band size: 22 kDa



Paraformaldehyde-fixed, paraffin embedded (rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (GPX2) Polyclonal Antibody, Unconjugated (AP55158) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.