

ZMYM2 Rabbit pAb

ZMYM2 Rabbit pAb

Catalog # AP55217

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9UBW7
Reactivity	Rat, Human
Predicted	Mouse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	154911
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ZMYM2/ZNF198
Epitope Specificity	1051-1150/1377
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
DISEASE	Note=A chromosomal aberration involving ZMYM2 may be a cause of stem cell leukemia lymphoma syndrome (SCLL). Translocation t(8;13)(p11;q12) with FGFR1. SCLL usually presents as lymphoblastic lymphoma in association with a myeloproliferative disorder, often accompanied by pronounced peripheral eosinophilia and/or prominent eosinophilic infiltrates in the affected bone marrow.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZNF198 (zinc finger 198), also known as ZMYM2 (zinc finger, MYM-type 2), FIM, MYM, RAMP or SCLL, is a 1,377 amino acid protein that localizes to the nucleus and contains nine MYM-type zinc fingers. Thought to be a component of the BHC histone deacetylase complex, ZNF198 interacts with HDAC1 and HDAC2 and is thought to stabilize the BHC complex via its MYM-type zinc fingers. The gene encoding ZNF198 is subject to a translocation with Flg, an event that may be involved in the pathogenesis of stem cell leukemia lymphoma syndrome (SCLL), a lymphoblastic lymphoma often accompanied by pronounced peripheral eosinophilia and/or prominent eosinophilic infiltrates in the affected bone marrow.

Additional Information

Gene ID	7750
Other Names	Zinc finger MYM-type protein 2, Fused in myeloproliferative disorders protein, Rearranged in atypical myeloproliferative disorder protein, Zinc finger protein

198, ZMYM2, FIM, RAMP, ZNF198

Dilution

IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name

ZMYM2

Synonyms

FIM, RAMP, ZNF198

Function

Involved in the negative regulation of transcription.


Cellular Location

Nucleus.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images




Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZMYM2) Polyclonal Antibody, Unconjugated (AP55217) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-ZMYM2 Polyclonal Antibody, Unconjugated (AP55217) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining

Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZMYM2) Polyclonal Antibody, Unconjugated (AP55217) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit)



(sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.