

CHST7 Rabbit pAb

CHST7 Rabbit pAb
Catalog # AP55362

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9NS84
Reactivity	Mouse
Predicted	Human, Rat, Dog, Pig, Rabbit, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54266
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CHST7
Epitope Specificity	101-200/486
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Golgi apparatus membrane; Single-pass type II membrane protein
SIMILARITY	Belongs to the sulfotransferase 1 family. Gal/GlcNAc/GalNAc subfamily.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic states of DNA during transcription. This enzyme catalyzes the transient breaking and rejoining of a single strand of DNA which allows the strands to pass through one another, thus reducing the number of supercoils and altering the topology of DNA. This enzyme forms a complex with BLM which functions in the regulation of recombination in somatic cells. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	56548
Other Names	Carbohydrate sulfotransferase 7, 2.8.2.-, 2.8.2.17, Chondroitin 6-sulfotransferase 2, C6ST-2, Galactose/N-acetylglucosamine/N-acetylglucosamine 6-O-sulfotransferase 5, GST-5, N-acetylglucosamine 6-O-sulfotransferase 4, GlcNAc6ST-4, Gn6st-4, CHST7
Target/Specificity	Widely expressed. Highly expressed in heart, spleen, liver and ovary. Expressed at lower level in brain, placenta, thyroid, spinal cord and peripheral blood leukocytes. Not expressed in adult skin.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information**Name**

CHST7

Function

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of non-reducing N-acetylglucosamine (GlcNAc) residues. Preferentially acts on mannose-linked GlcNAc. Also able to catalyze the transfer of sulfate to position 6 of the N-acetylgalactosamine (GalNAc) residue of chondroitin. Also acts on core 2 mucin-type oligosaccharide and N-acetyllactosamine oligomer with a lower efficiency. Has weak or no activity toward keratan sulfate and oligosaccharides containing the Galbeta1-4GlcNAc. Catalyzes 6-O-sulfation of beta-benzyl GlcNAc but not alpha- or beta-benzyl GalNAc.

Cellular Location


Golgi apparatus membrane; Single-pass type II membrane protein

Tissue Location

Widely expressed. Highly expressed in heart, spleen, liver and ovary. Expressed at lower level in brain, placenta, thyroid, spinal cord and peripheral blood leukocytes. Not expressed in adult skin.


Background

This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic states of DNA during transcription. This enzyme catalyzes the transient breaking and rejoining of a single strand of DNA which allows the strands to pass through one another, thus reducing the number of supercoils and altering the topology of DNA. This enzyme forms a complex with BLM which functions in the regulation of recombination in somatic cells. [provided by RefSeq, Jul 2008]

Images

Sample: Kidney (Mouse) Lysate at 40 ug Cerebrum (Mouse) Lysate at 40 ug Primary: Anti- CHST7 (AP55362) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 54 kD Observed band size: 54 kD

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CHST7) Polyclonal Antibody, Unconjugated (AP55362) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit)



(sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.