

# COQ3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55368

## Product Information

---

<b>Application</b>	IHC-P, IHC-F, IF, ICC, E
<b>Primary Accession</b>	<a href="#">Q9NZJ6</a>
<b>Reactivity</b>	Rat, Pig, Dog, Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	41054
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human COQ3
<b>Epitope Specificity</b>	151-250/369
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Mitochondrion matrix.
<b>SIMILARITY</b>	Belongs to the methyltransferase superfamily. UbiG/COQ3 family.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	Ubiquinone, also known as coenzyme Q, or Q, is a critical component of the electron transport pathways of both eukaryotes and prokaryotes (Jonassen and Clarke, 2000 [PubMed 10777520]). This lipid consists of a hydrophobic isoprenoid tail and a quinone head group. The tail varies in length depending on the organism, but its purpose is to anchor coenzyme Q to the membrane. The quinone head group is responsible for the activity of coenzyme Q in the respiratory chain. The <i>S. cerevisiae</i> COQ3 gene encodes an O-methyltransferase required for 2 steps in the biosynthetic pathway of coenzyme Q. This enzyme methylates an early coenzyme Q intermediate, 3,4-dihydroxy-5-polyprenylbenzoic acid, as well as the final intermediate in the pathway, converting demethyl-ubiquinone to coenzyme Q. The COQ3 gene product is also capable of methylating the distinct prokaryotic early intermediate 2-hydroxy-6-polyprenyl phenol.[supplied by OMIM, Mar 2008]

## Additional Information

---

<b>Gene ID</b>	51805
<b>Other Names</b>	Ubiquinone biosynthesis O-methyltransferase, mitochondrial {ECO:0000255 HAMAP-Rule:MF_03190}, 3-demethylubiquinol 3-O-methyltransferase {ECO:0000255 HAMAP-Rule:MF_03190}, 2.1.1.64 {ECO:0000255 HAMAP-Rule:MF_03190}, Polyprenyldihydroxybenzoate methyltransferase {ECO:0000255 HAMAP-Rule:MF_03190}, 2.1.1.114 {ECO:0000255 HAMAP-Rule:MF_03190}, COQ3 {ECO:0000255 HAMAP-Rule:MF_03190}

<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
<b>Format</b>	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

---

<b>Name</b>	COQ3 {ECO:0000255   HAMAP-Rule:MF_03190, ECO:0000303   PubMed:38425362}
<b>Function</b>	O-methyltransferase required for two non-consecutive steps during ubiquinone biosynthesis (By similarity) (PubMed: <a href="#">10777520</a> , PubMed: <a href="#">38425362</a> ). Catalyzes the 2 O-methylation of 3,4-dihydroxy-5-(all-trans-decaprenyl)benzoic acid into 4-hydroxy-3-methoxy-5-(all-trans-decaprenyl)benzoic acid (By similarity) (PubMed: <a href="#">10777520</a> , PubMed: <a href="#">38425362</a> ). Also catalyzes the last step of ubiquinone biosynthesis by mediating methylation of 3-demethylubiquinone into ubiquinone (By similarity) (PubMed: <a href="#">38425362</a> ). Also able to mediate the methylation of 3-demethylubiquinol-10 into ubiquinol-10 (By similarity) (PubMed: <a href="#">10777520</a> ).
<b>Cellular Location</b>	Mitochondrion inner membrane {ECO:0000255   HAMAP- Rule:MF_03190, ECO:0000269   PubMed:27499296}; Peripheral membrane protein {ECO:0000255   HAMAP-Rule:MF_03190}; Matrix side {ECO:0000255   HAMAP-Rule:MF_03190}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.