

EOLA1/CXorf40A Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55431

Product Information

| Application | IHC-P, IHC-F, IF, ICC, E |
|-------------------|--------------------------|
| Primary Accession | <u>Q8TE69</u> |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 17891 |

Additional Information

| Gene ID | 91966 |
|-------------|--|
| Other Names | Protein EOLA1, Endothelial-overexpressed lipopolysaccharide-associated factor 1, Endothelium and lymphocyte associated ASCH domain 1 {ECO:0000312 HGNC:HGNC:28089}, EOLA1 (<u>HGNC:28089</u>), CXorf40, CXorf40A |
| Dilution | IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000- 10000 |
| Format | 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce |
| Storage | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |

| Protein Information | |
|---------------------|---|
| Name | EOLA1 (<u>HGNC:28089</u>) |
| Synonyms | CXorf40, CXorf40A |
| Function | May play a role in cell protection during the inflammatory response. In epithelial cells, negatively regulates IL6 production and apoptosis through the regulation of MT2A expression (PubMed: <u>24916366</u>). |
| Tissue Location | Expressed primarily in heart, skeletal muscle, kidney, liver and placenta. Relatively high level of expression in spleen, colon and small intestine. Almost no expression in brain, thymus, lung and peripheral blood leukocytes. Expressed in epithelial cells (at protein level) (PubMed:31007603) |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.