

# NSG1/D4S234E Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55451

## Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">P42857</a>
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20913

## Additional Information

Gene ID	27065
Other Names	Neuronal vesicle trafficking-associated protein 1, Neuron-enriched endosomal protein of 21 kDa, Neuron-specific protein family member 1 {ECO:0000312 HGNC:HGNC:18790}, NSG1 ( <a href="#">HGNC:18790</a> )
Dilution	Elisa=1:500-1000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ICC=1:100-500,
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## Protein Information

Name	NSG1 ( <a href="#">HGNC:18790</a> )
Function	Plays a role in the recycling mechanism in neurons of multiple receptors, including AMPAR, APP and L1CAM and acts at the level of early endosomes to promote sorting of receptors toward a recycling pathway. Regulates sorting and recycling of GRIA2 through interaction with GRIP1 and then contributes to the regulation of synaptic transmission and plasticity by affecting the recycling and targeting of AMPA receptors to the synapse (By similarity). Is required for faithful sorting of L1CAM to axons by facilitating trafficking from somatodendritic early endosome or the recycling endosome (By similarity). In an other hand, induces apoptosis via the activation of CASP3 in response to DNA damage (PubMed: <a href="#">20599942</a> , PubMed: <a href="#">20878061</a> ).
Cellular Location	Membrane {ECO:0000250 UniProtKB:P02683}; Single- pass type II membrane protein {ECO:0000250 UniProtKB:P02683}. Golgi apparatus, trans-Golgi network membrane {ECO:0000250 UniProtKB:P02683} Endosome membrane {ECO:0000250 UniProtKB:P02683}. Cell projection, dendrite {ECO:0000250 UniProtKB:P02683}. Early endosome membrane

{ECO:0000250|UniProtKB:P02683}. Late endosome membrane  
{ECO:0000250|UniProtKB:P02683}. Lysosome lumen  
{ECO:0000250|UniProtKB:P02683}. Recycling endosome membrane  
{ECO:0000250|UniProtKB:P02683}. Cytoplasmic vesicle membrane  
{ECO:0000250|UniProtKB:P02683}. Golgi apparatus, Golgi stack membrane  
{ECO:0000250|UniProtKB:P02683}. Endosome, multivesicular body  
membrane {ECO:0000250|UniProtKB:P02683}. Endoplasmic reticulum  
membrane. Note=Endocytosed from the cell surface, thus enters into early  
endosomes, trafficks to late endosomes and degradates in lysosomes (By  
similarity). Endoplasmic reticulum targeting is essential for apoptosis  
(PubMed:20599942). Found in both stationary and motile endosomes. A  
previous study supports a type I membrane protein topology (By similarity)  
{ECO:0000250|UniProtKB:P02683, ECO:0000250|UniProtKB:Q62092,  
ECO:0000269|PubMed:20599942}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.