

DDX26/DICE1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55469

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9UL03
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	100390

Additional Information

Gene ID	26512
Other Names	Integrator complex subunit 6, Int6, DBI-1, Protein DDX26, Protein deleted in cancer 1, DICE1, INTS6, DBI1, DDX26, DDX26A
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	INTS6 {ECO:0000303 PubMed:33243860, ECO:0000312 HGNC:HGNC:14879}
Function	Component of the integrator complex, a multiprotein complex that terminates RNA polymerase II (Pol II) transcription in the promoter-proximal region of genes (PubMed:33243860, PubMed:34004147, PubMed:39504960). The integrator complex provides a quality checkpoint during transcription elongation by driving premature transcription termination of transcripts that are unfavorably configured for transcriptional elongation: the complex terminates transcription by (1) catalyzing dephosphorylation of the C-terminal domain (CTD) of Pol II subunit POLR2A and SUPT5H/SPT5, (2) degrading the exiting nascent RNA transcript via endonuclease activity and (3) promoting the release of Pol II from bound DNA (PubMed:33243860, PubMed:34004147, PubMed:38570683, PubMed:39504960). The integrator complex is also involved in terminating the synthesis of non-coding Pol II transcripts, such as enhancer RNAs (eRNAs), small nuclear RNAs (snRNAs), telomerase RNAs and long non-coding RNAs (lncRNAs) (PubMed:16239144). Within the integrator complex, INTS6 acts as a molecular adapter that promotes assembly of protein phosphatase 2A (PP2A) subunits to the integrator core complex, promoting recruitment of PP2A to transcription pause- release checkpoint

(PubMed:[33243860](#), PubMed:[34004147](#)). Mediates recruitment of cytoplasmic dynein to the nuclear envelope, probably as component of the integrator complex (PubMed:[23904267](#)). May have a tumor suppressor role; an ectopic expression suppressing tumor cell growth (PubMed:[15254679](#), PubMed:[16239144](#)).

Cellular Location

Nucleus. Chromosome Note=Associates with chromatin and transcription pause-release checkpoint.

Tissue Location

Widely expressed. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.