

DEFB119 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55483

Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession

Reactivity

Dog

Host

Clonality

Polyclonal

Calculated MW

Physical State

Q8N690

Pog

Rabbit

Polyclonal

9822

Liquid

Immunogen KLH conjugated synthetic peptide derived from human DEFB119

Epitope Specificity 22-84/84 Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Secreted.

SIMILARITY Belongs to the beta-defensin family.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions This gene encodes a member of a family of small secreted proteins. These

proteins participate in immune defense against microbial infection. This gene is located in a cluster of similar genes on chromosome 20. Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Sep 2012]

Additional Information

Gene ID 245932

Other Names Beta-defensin 119, Beta-defensin 120, Beta-defensin 19, DEFB-19,

Beta-defensin 20, DEFB-20, Defensin, beta 119, Defensin, beta 120,

ESC42-RELA, DEFB119, DEFB120, DEFB19, DEFB20

Dilution IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-

10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name DEFB119

Synonyms DEFB120, DEFB19, DEFB20

Has antibacterial activity. **Function**

Cellular Location Secreted.

Tissue Location

Abundant expression in the male reproductive tract only. Abundant expressed in testis and the caput region of epididymis, but low in the corpus

region.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.