

DEFB121 Rabbit pAb

DEFB121 Rabbit pAb

Catalog # AP55484

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q8NES8
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	8058
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DEFB121
Epitope Specificity	16-76/76
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Beta-defensins are small cationic peptides with broad-spectrum antimicrobial activity against a variety of enveloped viruses, fungi and bacteria. Produced in mucosal epithelia and neutrophils of several species, Beta-defensins are developmentally regulated. The family of Beta-defensin proteins share a common defensin-motif that is characterized by multiple cysteine residues and a highly conserved tertiary structure. Besides playing a significant role in host immune defense, many Beta-defensins also are involved in sperm maturation and capacitation. Beta-defensin 124 is a 71 amino acid secreted protein that most likely contains a signal peptide sequence that requires cleavage by proteolytic enzymes in order to become biologically active.

Additional Information

Gene ID	245937
Other Names	Beta-defensin 124, Beta-defensin 24, DEFB-24, Defensin, beta 124, DEFB124, DEFB24
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	DEFB124
Synonyms	DEFB24
Function	Has antibacterial activity.
Cellular Location	Secreted.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DEFB121) Polyclonal Antibody, Unconjugated (AP55484) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.