

DEPDC1B Rabbit pAb

DEPDC1B Rabbit pAb

Catalog # AP55500

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q8WUY9
Reactivity	Human
Predicted	Rat, Mouse, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61771
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DEPDC1B
Epitope Specificity	451-529/529
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	intracellular
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	GTPase activator activity. biological process: intracellular signaling cascade

Additional Information

Gene ID	55789
Other Names	DEP domain-containing protein 1B, HBV X-transactivated gene 8 protein, HBV XAg-transactivated protein 8, DEPDC1B, XTP8
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	DEPDC1B
Synonyms	XTP8

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:

Lane 1: Human HepG2 cell lysates


Lane 2: Human MCF-7 cell lysates

Primary: Anti-DEPDC1B/BRCC3 (AP55500) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 62 kDa

Observed band size: 68 kDa



Tissue/cell: human brain glioma; 4%

Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-BRCC3 Polyclonal Antibody, Unconjugated(AP55500) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.