

DHX33 Rabbit pAb

DHX33 Rabbit pAb
Catalog # AP55520

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9H6R0
Reactivity	Mouse
Predicted	Rat, Dog, Human, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78874
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DHX33
Epitope Specificity	101-200/707
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus, nucleolus. Nucleus, nucleoplasm. Note=Predominantly in the nucleolus. During mitosis, localizes with the nucleolar organizing regions.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the DEAD box protein family. The DEAD box proteins are characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this DEAD box protein family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2010]

Additional Information

Gene ID	56919
Other Names	ATP-dependent RNA helicase DHX33, 3.6.4.13, DEAH box protein 33, DHX33 (HGNC:16718), DDX33
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	DHX33 (HGNC:16718)
Synonyms	DDX33
Function	Implicated in nucleolar organization, ribosome biogenesis, protein synthesis and cytoplasmic dsRNA sensing (By similarity) (PubMed: 21930779 , PubMed: 23871209 , PubMed: 26100019). Stimulates RNA polymerase I transcription of the 47S precursor rRNA. Associates with ribosomal DNA (rDNA) loci where it is involved in POLR1A recruitment (PubMed: 21930779). In the cytoplasm, promotes elongation-competent 80S ribosome assembly at the late stage of mRNA translation initiation (PubMed: 26100019). Senses cytosolic dsRNA mediating NLRP3 inflammasome formation in macrophages and type I interferon production in myeloid dendritic cells (PubMed: 23871209). Required for NLRP3 activation induced by viral dsRNA and bacterial RNA (PubMed: 23871209). In dendritic cells, required for induction of type I interferon production induced by cytoplasmic dsRNA via the activation of MAPK and NF-kappa-B signaling pathways (By similarity).
Cellular Location	Nucleus, nucleolus. Nucleus, nucleoplasm. Cytoplasm. Nucleus Inflammasome. Note=Predominantly in the nucleolus. During mitosis, localizes with the nucleolar organizing regions (PubMed:21930779). Upon dsRNA-binding, localizes in the inflammasome (PubMed:23871209).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DHX33) Polyclonal Antibody, Unconjugated (AP55520) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.