

DIMT1L Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55525

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9UNQ2
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35236

Additional Information

Gene ID	27292
Other Names	Probable dimethyladenosine transferase, 2.1.1.183, DIM1 dimethyladenosine transferase 1 homolog, DIM1 dimethyladenosine transferase 1-like, Probable 18S rRNA (adenine(1779)-N(6)/adenine(1780)-N(6))-dimethyltransferase, Probable 18S rRNA dimethylase, Probable S-adenosylmethionine-6-N', N'-adenosyl(rRNA) dimethyltransferase, DIMT1, DIMT1L
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DIMT1 (HGNC:30217)
Synonyms	DIMT1L
Function	Specifically dimethylates two adjacent adenosines in the loop of a conserved hairpin near the 3'-end of 18S rRNA in the 40S particle (PubMed: 25851604). Involved in the pre-rRNA processing steps leading to small-subunit rRNA production independently of its RNA-modifying catalytic activity (PubMed: 25851604). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed: 34516797).

Cellular Location

Nucleus, nucleoplasm. Nucleus, nucleolus

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.