

EHBP1 Rabbit pAb

EHBP1 Rabbit pAb
Catalog # AP55610

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q8NDI1
Reactivity	Rat, Human
Predicted	Pig, Mouse, Rabbit, Dog, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	140017
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human EHBP1
Epitope Specificity	21-120/1231
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell Membrane and Cytoplasmic
DISEASE	Prostate cancer, hereditary, 12 (HPC12) [MIM:611868]: A condition associated with familial predisposition to cancer of the prostate. Most prostate cancers are adenocarcinomas that develop in the acini of the prostatic ducts. Other rare histopathologic types of prostate cancer that occur in approximately 5% of patients include small cell carcinoma, mucinous carcinoma, prostatic ductal carcinoma, transitional cell carcinoma, squamous cell carcinoma, basal cell carcinoma, adenoid cystic carcinoma (basaloid), signet-ring cell carcinoma and neuroendocrine carcinoma. Note=Disease susceptibility is associated with variations affecting the gene represented in this entry.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes an Eps15 homology domain binding protein. The encoded protein may play a role in endocytic trafficking. A single nucleotide polymorphism in this gene is associated with an aggressive form of prostate cancer. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Feb 2010]

Additional Information

Gene ID	23301
Other Names	EH domain-binding protein 1, EHBP1, KIAA0903, NACSIN
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.


Protein Information

Name	EHBP1
Synonyms	KIAA0903, NACSIN
Function	May play a role in actin reorganization. Links clathrin- mediated endocytosis to the actin cytoskeleton. May act as Rab effector protein and play a role in vesicle trafficking (PubMed: 14676205 , PubMed: 27552051). Required for perinuclear sorting and insulin- regulated recycling of SLC2A4/GLUT4 in adipocytes (By similarity).
Cellular Location	Cytoplasm. Membrane. Endosome Note=Mostly found in cytosol and plasma membrane


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample:
293T(Human) Cell Lysate at 40 ug
Primary: Anti-EHBP1 (AP55610) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 140 kD
Observed band size: 134 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain);
Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (EHBP1) Polyclonal Antibody, Unconjugated (AP55610) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.