

EIF2B2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55615

Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	P49770
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38990

Additional Information

Gene ID	8892
Other Names	Translation initiation factor eIF-2B subunit beta, S20I15, S20III15, eIF-2B GDP-GTP exchange factor subunit beta, EIF2B2, EIF2BB
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	EIF2B2
Synonyms	EIF2BB
Function	Acts as a component of the translation initiation factor 2B (eIF2B) complex, which catalyzes the exchange of GDP for GTP on eukaryotic initiation factor 2 (eIF2) gamma subunit (PubMed: 25858979 , PubMed: 27023709 , PubMed: 31048492). Its guanine nucleotide exchange factor activity is repressed when bound to eIF2 complex phosphorylated on the alpha subunit, thereby limiting the amount of methionyl- initiator methionine tRNA available to the ribosome and consequently global translation is repressed (PubMed: 25858979 , PubMed: 31048492).
Cellular Location	Cytoplasm, cytosol {ECO:0000250 UniProtKB:Q9UT76}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.