

eIF4A3 Rabbit pAb

eIF4A3 Rabbit pAb Catalog # AP55620

Product Information

Application WB, IHC-P, IHC-F, IF

Primary Accession <u>P38919</u>

Reactivity Human, Mouse, Dog

Host Rabbit
Clonality Polyclonal
Calculated MW 46871
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human eIF4A3

Epitope Specificity 351-411/411

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus. Nucleus speckle. Cytoplasm. Nucleocytoplasmic shuttling protein.

Travels to the cytoplasm as part of the exon junction complex (EJC) bound to mRNA. Detected in dendritic layer as well as the nuclear and cytoplasmic (somatic) compartments of neurons. Colocalizes with STAU1 and FMR1 in

dendrites.

SIMILARITY Belongs to the DEAD box helicase family. eIF4A subfamily. Contains 1 helicase

ATP-binding domain. Contains 1 helicase C-terminal domain.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions This gene encodes a member of the DEAD box protein family. DEAD box

proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene is a nuclear matrix

protein. Its amino acid sequence is highly similar to the amino acid sequences of the translation initiation factors eIF4AI and eIF4AII, two other members of

the DEAD box protein family. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID 9775

Other Names Eukaryotic initiation factor 4A-III, eIF-4A-III, eIF4A-III, 3.6.4.13,

ATP-dependent RNA helicase DDX48, ATP-dependent RNA helicase eIF4A-3, DEAD box protein 48, Eukaryotic initiation factor 4A-like NUK-34, Eukaryotic translation initiation factor 4A isoform 3, Nuclear matrix protein 265, NMP

265, hNMP 265, Eukaryotic initiation factor 4A-III, N-terminally processed,

EIF4A3, DDX48, KIAA0111

Target/Specificity Ubiquitously expressed.

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-

500,Flow-Cyt=2ug/Test

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name EIF4A3

Synonyms DDX48, KIAA0111

Function ATP-dependent RNA helicase (PubMed: <u>16170325</u>). Involved in pre-mRNA

splicing as component of the spliceosome (PubMed: 11991638, PubMed: 22961380, PubMed: 28076346, PubMed: 28502770,

PubMed: <u>29301961</u>). Core component of the splicing-dependent multiprotein

exon junction complex (EJC) deposited at splice junctions on mRNAs

(PubMed:<u>16170325</u>, PubMed:<u>16209946</u>, PubMed:<u>16314458</u>, PubMed:<u>16923391</u>, PubMed:<u>16931718</u>, PubMed:<u>19033377</u>,

PubMed: 20479275). The EJC is a dynamic structure consisting of core proteins and several peripheral nuclear and cytoplasmic associated factors that join the complex only transiently either during EJC assembly or during subsequent

mRNA metabolism. The EJC marks the position of the exon-exon junction in the mature mRNA for the gene expression machinery and the core components remain bound to spliced mRNAs throughout all stages of mRNA metabolism thereby influencing downstream processes including nuclear mRNA export, subcellular mRNA localization, translation efficiency and nonsense- mediated mRNA decay (NMD). Its RNA-dependent ATPase and RNA-helicase activities are induced by CASC3, but abolished in presence of the MAGOH-RBM8A heterodimer, thereby trapping the ATP-bound EJC core onto spliced mRNA in a stable conformation. The inhibition of ATPase activity by the MAGOH-RBM8A heterodimer increases the RNA-binding affinity of the EJC. Involved in translational enhancement of spliced mRNAs after formation of the 80S ribosome complex. Binds spliced mRNA in sequence-independent manner, 20-24 nucleotides upstream of mRNA exon-exon junctions. Shows higher affinity for single-stranded RNA in an ATP-bound core EJC complex than after the ATP is hydrolyzed. Involved in the splicing modulation of BCL2L1/Bcl-X (and probably other apoptotic genes); specifically inhibits formation of proapoptotic isoforms such as Bcl-X(S); the function is different

from the established EJC assembly (PubMed: <u>22203037</u>). Involved in craniofacial development (PubMed: <u>24360810</u>).

Cellular Location Nucleus. Nucleus speckle. Cytoplasm {ECO:0000250|UniProtKB:Q3B8Q2}.

Note=Nucleocytoplasmic shuttling protein. Travels to the cytoplasm as part of the exon junction complex (EJC) bound to mRNA. Detected in dendritic layer as well as the nuclear and cytoplasmic (somatic) compartments of neurons.

Colocalizes with STAU1 and FMR1 in dendrites (By similarity)

{ECO:0000250|UniProtKB:Q3B8Q2}


Tissue Location Ubiquitously expressed.

2 of 4

Background

This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene is a nuclear matrix protein. Its amino acid sequence is highly similar to the amino acid sequences of the translation initiation factors eIF4AI and eIF4AII, two other members of the DEAD box protein family. [provided by RefSeq, Jul 2008]

Images



Blank control:Hela.

Primary Antibody (green line): Rabbit Anti-eIF4A3


antibody (AP55620) Dilution: 2ug/Test;

Secondary Antibody: Goat anti-rabbit IgG-FITC


Dilution: 0.5ug/Test.

Protocol

The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature . Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (eIF4A3) polyclonal Antibody, Unconjugated (AP55620) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Sample:

Lane 1: Mouse Testis tissue lysates

Lane 2: Rat Testis tissue lysates

Lane 3: Human HepG2 cell lysates

Lane 4: Human HeLa cell lysates

Lane 5: Human MCF-7 cell lysates


Primary: Anti-eIF4A3 (AP55620) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000

dilution

Predicted band size: 47 kDa Observed band size: 47 kDa

Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate



buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (eIF4A3) Polyclonal Antibody, Unconjugated (AP55620) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.