

ELP4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55626

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q96EB1
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46588
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ELP4
Epitope Specificity	321-424/424
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Nucleus.
SIMILARITY	Belongs to the ELP4 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a membrane-bound protein which is a member of the ELO family, proteins which participate in the biosynthesis of fatty acids. Consistent with the expression of the encoded protein in photoreceptor cells of the retina, mutations and small deletions in this gene are associated with Stargardt-like macular dystrophy (STGD3) and autosomal dominant Stargardt-like macular dystrophy (ADMD), also referred to as autosomal dominant atrophic macular degeneration. [provided by RefSeq, Jul 2008]


Additional Information

Gene ID	26610
Other Names	Elongator complex protein 4, hELP4, PAX6 neighbor gene protein, ELP4, C11orf19, PAXNEB
Target/Specificity	Widely expressed
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	ELP4
Synonyms	C11orf19, PAXNEB
Function	Component of the elongator complex which is required for multiple tRNA modifications, including mcm5U (5-methoxycarbonylmethyl uridine), mcm5s2U (5-methoxycarbonylmethyl-2-thiouridine), and ncm5U (5-carbamoylmethyl uridine) (PubMed: 29332244). The elongator complex catalyzes the formation of carboxymethyluridine in the wobble base at position 34 in tRNAs (PubMed: 29332244).
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Widely expressed..


Images



Protein: HeLa(human) lysate at 40ug;
Primary: rabbit Anti-ELP4 (AP55626) at 1:300;
Secondary: HRP conjugated Goat-Anti-rabbit
IgG(bs-0295G-HRP) at 1: 5000;
Predicted band size: 47 kD
Observed band size: 47 kD




Paraformaldehyde-fixed, paraffin embedded (rat brain);
Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ELP4) Polyclonal Antibody, Unconjugated (AP55626) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ELP4) Polyclonal Antibody, Unconjugated (AP55626) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase



by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ELP4) Polyclonal Antibody, Unconjugated (AP55626) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.