

EMC9 Rabbit pAb

EMC9 Rabbit pAb
Catalog # AP55629

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9Y3B6
Predicted	Rat, Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23061
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human EMC9
Epitope Specificity	151-250/208
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasmic
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Component of the ER membrane protein complex (EMC).

Additional Information

Gene ID	51016
Other Names	ER membrane protein complex subunit 9, Protein FAM158A, EMC9, C14orf122, FAM158A
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	EMC9
Synonyms	C14orf122, FAM158A
Function	Part of the endoplasmic reticulum membrane protein complex (EMC) that enables the energy-independent insertion into endoplasmic reticulum

membranes of newly synthesized membrane proteins (PubMed:[29242231](#), PubMed:[29809151](#), PubMed:[30415835](#), PubMed:[32459176](#)). Preferentially accommodates proteins with transmembrane domains that are weakly hydrophobic or contain destabilizing features such as charged and aromatic residues (PubMed:[29242231](#), PubMed:[29809151](#), PubMed:[30415835](#)). Involved in the cotranslational insertion of multi-pass membrane proteins in which stop-transfer membrane-anchor sequences become ER membrane spanning helices (PubMed:[29809151](#), PubMed:[30415835](#)). It is also required for the post-translational insertion of tail-anchored/TA proteins in endoplasmic reticulum membranes (PubMed:[29242231](#), PubMed:[29809151](#)). By mediating the proper cotranslational insertion of N-terminal transmembrane domains in an N-exo topology, with translocated N-terminus in the lumen of the ER, controls the topology of multi-pass membrane proteins like the G protein-coupled receptors (PubMed:[30415835](#)). By regulating the insertion of various proteins in membranes, it is indirectly involved in many cellular processes (Probable).

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.