

FAM175B Rabbit pAb

FAM175B Rabbit pAb

Catalog # AP55707

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q15018
Reactivity	Mouse
Predicted	Rat, Pig, Human, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46901
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FAM175B
Epitope Specificity	41-140/415
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	FAM175B (ABRO1) is a 415 amino acid protein, sharing 39% identity to ABRA1 in the N terminal two thirds of the protein (amino acids 1 to 260), including a coiled-coil domain and a region called the ABR domain. However, ABRO1 lacks the phosphorylated ser-X-X-phe motif of ABRA1 and, unlike ABRA1, does not bind to BRCA1.

Additional Information

Gene ID	23172
Other Names	BRISC complex subunit Abraxas 2 {ECO:0000312 HGNC:HGNC:28975}, Abraxas brother protein 1, Protein FAM175B, ABRAXAS2 (HGNC:28975)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ABRAXAS2 (HGNC:28975)
Function	Component of the BRISC complex, a multiprotein complex that specifically

cleaves 'Lys-63'-linked polyubiquitin, leaving the last ubiquitin chain attached to its substrates (PubMed:[19214193](#), PubMed:[20032457](#), PubMed:[20656690](#), PubMed:[24075985](#)). May act as a central scaffold protein that assembles the various components of the BRISC complex and retains them in the cytoplasm (PubMed:[20656690](#)). Plays a role in regulating the onset of apoptosis via its role in modulating 'Lys-63'-linked ubiquitination of target proteins (By similarity). Required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1 (PubMed:[26195665](#)). Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activities by enhancing its stability and cell surface expression (PubMed:[24075985](#), PubMed:[26344097](#)). Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination (PubMed:[24075985](#)). Required for normal induction of p53/TP53 in response to DNA damage (PubMed:[25283148](#)). Independent of the BRISC complex, promotes interaction between USP7 and p53/TP53, and thereby promotes deubiquitination of p53/TP53, preventing its degradation and resulting in increased p53/TP53-mediated transcription regulation and p53/TP53-dependent apoptosis in response to DNA damage (PubMed:[25283148](#)).

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton. Note=A minor proportion is detected in the nucleus (PubMed:21282113, PubMed:22974638). Translocates into the nucleus in response to DNA damage (PubMed:25283148). Directly binds to microtubules and is detected at the minus end of K-fibers (PubMed:26195665). Co-localizes with NUMA1 at mitotic spindle poles (PubMed:26195665).


Tissue Location

Detected in heart muscle (at protein level). Detected in heart and muscle, and at much lower levels in brain (PubMed:21195082).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM175B) Polyclonal Antibody, Unconjugated (AP55707) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.