

C1orf156 Rabbit pAb

C1orf156 Rabbit pAb

Catalog # AP55749

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	O95568
Reactivity	Human
Predicted	Rat, Pig, Dog, Mouse, Rabbit, Chicken, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42148
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C1orf156
Epitope Specificity	121-220/372
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The C1orf156 gene product has been provisionally designated C1orf156 pending further characterization.

Additional Information

Gene ID	92342
Other Names	Histidine protein methyltransferase 1 homolog, 2.1.1.85, Arsenic-transactivated protein 2 {ECO:0000303 Ref.1}, AsTP2 {ECO:0000303 Ref.1}, Methyltransferase-like protein 18 {ECO:0000312 HGNC:HGNC:28793}, METTL18 {ECO:0000303 PubMed:33693809, ECO:0000312 HGNC:HGNC:28793}

Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	METTL18 {ECO:0000303 PubMed:33693809, ECO:0000312 HGNC:HGNC:28793}
Function	Protein-L-histidine N-tele-methyltransferase that specifically monomethylates RPL3, thereby regulating translation elongation (PubMed: 23349634 , PubMed: 33693809 , PubMed: 35674491). Histidine methylation of RPL3 regulates translation elongation by slowing ribosome traversal on tyrosine codons: slower elongation provides enough time for proper folding of synthesized proteins and prevents cellular aggregation of tyrosine-rich proteins (PubMed: 35674491).
Cellular Location	Cytoplasm, cytosol. Nucleus. Nucleus, nucleolus

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C1orf156) Polyclonal Antibody, Unconjugated (AP55749) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.