

# CXCL14 Rabbit pAb

CXCL14 Rabbit pAb

Catalog # AP55754

## Product Information

---

<b>Application</b>	IHC-P, IHC-F, IF
<b>Primary Accession</b>	<a href="#">O95715</a>
<b>Reactivity</b>	Human, Rat
<b>Predicted</b>	Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	13078
<b>Physical State</b>	Liquid
<b>Immunogen</b>	KLH conjugated synthetic peptide derived from human CXCL14
<b>Epitope Specificity</b>	51-99/111
<b>Isotype</b>	IgG
<b>Purity</b>	affinity purified by Protein A
<b>Buffer</b>	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
<b>SUBCELLULAR LOCATION</b>	Secreted.
<b>SIMILARITY</b>	Belongs to the intercrine alpha (chemokine CxC) family.
<b>Important Note</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>Background Descriptions</b>	This gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than in inflammation. [provided by RefSeq, Jul 2008].

## Additional Information

---

<b>Gene ID</b>	9547
<b>Other Names</b>	C-X-C motif chemokine 14, Chemokine BRAK, MIP-2G, Small-inducible cytokine B14, CXCL14, MIP2G, NJAC, SCYB14
<b>Target/Specificity</b>	Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Highly expressed in normal tissue without inflammatory stimuli and infrequently expressed in cancer cell lines. Weakly expressed in monocyte-derived dendritic cells. Not detected in lung or unstimulated peripheral blood lymphocytes.
<b>Dilution</b>	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---


## Protein Information

<b>Name</b>	CXCL14
<b>Synonyms</b>	MIP2G, NJAC, SCYB14
<b>Function</b>	Potent chemoattractant for neutrophils, and weaker for dendritic cells. Not chemotactic for T-cells, B-cells, monocytes, natural killer cells or granulocytes. Does not inhibit proliferation of myeloid progenitors in colony formation assays.
<b>Cellular Location</b>	Secreted.
<b>Tissue Location</b>	Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Highly expressed in normal tissue without inflammatory stimuli and infrequently expressed in cancer cell lines. Weakly expressed in monocyte-derived dendritic cells. Not detected in lung or unstimulated peripheral blood lymphocytes

## Background

This gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than in inflammation. [provided by RefSeq, Jul 2008].

## Images



Tissue/cell: human lung carcinoma; 4%  
 Paraformaldehyde-fixed and paraffin-embedded;  
 Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling  
 bathing for 15min; Block endogenous peroxidase by 3%  
 Hydrogen peroxide for 30min; Blocking buffer (normal  
 goat serum,C-0005) at 37°C for 20 min;  
 Incubation: Anti-CXCL14 Polyclonal Antibody,  
 Unconjugated(AP55754) 1:200, overnight at 4°C, followed  
 by conjugation to the secondary antibody(SP-0023) and  
 DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.