

C1orf64 Rabbit pAb

C1orf64 Rabbit pAb

Catalog # AP55777

Product Information

Application	WB
Primary Accession	Q8NEQ6
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17657
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C1orf64
Epitope Specificity	81-169/169
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The C1orf64 gene product has been provisionally designated C1orf64 pending further characterization.

Additional Information

Gene ID	149563
Other Names	Steroid receptor-associated and regulated protein {ECO:0000312 HGNC:HGNC:28339}, Estrogen receptor-related factor, ER-related factor, Steroid receptor-regulated protein, SRARP (HGNC:28339)
Dilution	WB=1:500-2000

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	SRARP (HGNC:28339)
Function	May regulate the transcriptional function of androgen and estrogen receptors.
Tissue Location	Expressed in breast tumors with a higher expression level in estrogen receptor-positive cancers (PubMed:22341523, PubMed:28915724).

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Sample: HepG2 (human) cell Lysate at 40 ug
Primary: Anti- C1orf64 (AP55777) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 18 kD
Observed band size: 18 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.