10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

C3ORF31 Rabbit pAb

C3ORF31 Rabbit pAb
Catalog # AP55848

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q96BW9
Predicted	Rat, Pig, Dog, Human, Mouse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51067
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C3ORF31
Epitope Specificity	101-200/316
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Mitochondrion (By similarity).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	C3orf31 (chromosome 3 open reading frame 31), also known as MGC16471 or DKFZp434E0519, is a 316 amino acid mitochondrial protein that belongs to the MMP37 family and may be involved in translocation of transit peptide-containing proteins across the mitochondrial inner membrane. C3orf24 is encoded by a gene that maps to human chromosome 3p25.2. Chromosome 3 is made up of approximately 214 million bases encoding over 1,100 genes. Notably, there is a chemokine receptor gene cluster and a variety of human cancer related loci on chromosome 3. Particular regions of the chromosome 3 short arm are deleted in many types of cancer cells. Key tumor suppressing genes on chromosome 3 encode apoptosis mediator RASSF1, cell migration regulator HYAL1 and angiogenesis suppressor SEMA3B. Marfan Syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with chromosome 3.

Additional Information

Gene ID	132001
Other Names	Phosphatidate cytidylyltransferase, mitochondrial, 2.7.7.41, CDP-diacylglycerol synthase, CDP-DAG synthase, Mitochondrial translocator assembly and maintenance protein 41 homolog, TAM41, TAMM41, C3orf31
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TAMM41
Synonyms	C3orf31
Function	Catalyzes the conversion of phosphatidic acid (PA) to CDP- diacylglycerol (CDP-DAG), an essential intermediate in the synthesis of phosphatidylglycerol, cardiolipin and phosphatidylinositol.
Cellular Location	Mitochondrion inner membrane {ECO:0000250|UniProtKB:D3ZKT0}; Peripheral membrane protein {ECO:0000250|UniProtKB:D3ZKT0}; Matrix side {ECO:0000250|UniProtKB:P53230}

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.