

C3Orf34 Rabbit pAb

C3Orf34 Rabbit pAb

Catalog # AP55850

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q96LK0
Reactivity	Mouse
Predicted	Rat, Pig, Dog, Human, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19166
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C3Orf34
Epitope Specificity	81-163/163
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm, cytoskeleton, centrosome, centriole. Cytoplasm, cytoskeleton, spindle pole. Note=Associates with the mother centriole in early interphase. Localizes to spindle poles during mitosis, and to distinct foci oriented.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	C3orf34 (chromosome 3 open reading frame 34), also known as MGC14126, is a 163 amino acid protein encoded by a gene that maps to human chromosome 3q29. Chromosome 3 is made up of approximately 214 million bases encoding over 1,100 genes. Notably, there is a chemokine receptor gene cluster and a variety of human cancer related loci on chromosome 3. Particular regions of the chromosome 3 short arm are deleted in many types of cancer cells. Key tumor suppressing genes on chromosome 3 encode apoptosis mediator RASSF1, cell migration regulator HYAL1 and angiogenesis suppressor SEMA3B. Marfan Syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with chromosome 3.

Additional Information

Gene ID	84984
Other Names	Centrosomal protein of 19 kDa, Cep19, CEP19, C3orf34
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


Protein Information

Name	CEP19
Synonyms	C3orf34
Function	Required for ciliation (PubMed: 28428259 , PubMed: 28625565 , PubMed: 28659385). Recruits the RABL2B GTPase to the ciliary base to initiate ciliation. After specifically capturing the activated GTP- bound RABL2B, the CEP19-RABL2B complex binds intraflagellar transport (IFT) complex B from the large pool pre-docked at the base of the cilium and thus triggers its entry into the cilia (PubMed: 28428259 , PubMed: 28625565). Involved in the early steps in cilia formation by recruiting the ciliary vesicles (CVs) to the distal end of the mother centriole where they fuse to initiate cilium assembly. Involved in microtubule (MT) anchoring to the centrosomes (PubMed: 28659385).
Cellular Location	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, cilium basal body Note=Associates with the mother centriole in early interphase Localizes to spindle poles during mitosis, and to distinct foci oriented towards the midbody at telophase (PubMed:21399614). Localizes slightly apical to the subdistal appendage on the mother centriole, but below the distal appendage (PubMed:28625565, PubMed:28659385)

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C3Orf34) Polyclonal Antibody, Unconjugated (AP55850) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.