

C6ORF173 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55883

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q5EE01
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	10061
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C6ORF173
Epitope Specificity	31-88/88
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Chromosome, centromere. Chromosome, centromere, kinetochore. Nucleus matrix. Nucleus, nucleolus. Note=Constitutively localizes to centromeres throughout the cell cycle, and to the inner kinetochore during mitosis.
SIMILARITY	Belongs to the CENPW family.
SUBUNIT	Part of a centromere complex consisting of CENPA, CENPT and CENPW. Part of a centromere complex consisting of histone H3, CENPT and CENPW. Interacts directly with CENPT. Component of a heterotetrameric CENP-T-W-S-X complex composed of APITD1/CENPS, STRA13/CENPX, CENPT and CENPW. Interacts with NPM1. Binds DNA.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Component of the CENPA-NAC (nucleosome-associated) complex, a complex that plays a central role in assembly of kinetochore proteins, mitotic progression and chromosome segregation (By similarity). The CENPA-NAC complex recruits the CENPA-CAD (nucleosome distal) complex and may be involved in incorporation of newly synthesized CENPA into centromeres (By similarity). Part of a nucleosome-associated complex that binds specifically to histone H3-containing nucleosomes at the centromere, as opposed to nucleosomes containing CENPA. Component of the heterotetrameric CENP-T-W-S-X complex that binds and supercoils DNA, and plays an important role in kinetochore assembly. CENPW has a fundamental role in kinetochore assembly and function. It is one of the inner kinetochore proteins, with most further proteins binding downstream. Required for normal chromosome organization and normal progress through mitosis.

Additional Information

Gene ID 387103

Other Names	Centromere protein W, CENP-W, Cancer-up-regulated gene 2 protein, CENPW, C6orf173, CUG2
Target/Specificity	Highly expressed in ovary, liver, lung and pancreas and to a lower extent in breast and gastrointestinal tract cancers; such as those of the colon, rectum and stomach. Overexpressed in high grade breast invasive tumors. Expressed in many cancer cell types.
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CENPW
Synonyms	C6orf173, CUG2
Function	Component of the CENPA-NAC (nucleosome-associated) complex, a complex that plays a central role in assembly of kinetochore proteins, mitotic progression and chromosome segregation (By similarity). The CENPA-NAC complex recruits the CENPA-CAD (nucleosome distal) complex and may be involved in incorporation of newly synthesized CENPA into centromeres (By similarity). Part of a nucleosome-associated complex that binds specifically to histone H3-containing nucleosomes at the centromere, as opposed to nucleosomes containing CENPA. Component of the heterotetrameric CENP-T-W-S-X complex that binds and supercoils DNA, and plays an important role in kinetochore assembly. CENPW has a fundamental role in kinetochore assembly and function. It is one of the inner kinetochore proteins, with most further proteins binding downstream. Required for normal chromosome organization and normal progress through mitosis.
Cellular Location	Nucleus. Chromosome, centromere. Chromosome, centromere, kinetochore. Nucleus matrix. Nucleus, nucleolus. Note=Constitutively localizes to centromeres throughout the cell cycle, and to the inner kinetochore during mitosis. {ECO:0000250 UniProtKB:P0DJH6}
Tissue Location	Highly expressed in ovary, liver, lung and pancreas and to a lower extent in breast and gastrointestinal tract cancers; such as those of the colon, rectum and stomach. Overexpressed in high grade breast invasive tumors. Expressed in many cancer cell types

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.