

C9orf78 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP55957

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9NZ63
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33688
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human C9orf78
Epitope Specificity	21-120/289
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Chromosome 9 consists of about 145 million bases and 4% of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster. Chromosome 9 is partnered with chromosome 22 in the translocation leading to the aberrant production of BCR-ABL fusion protein often found in leukemias. The C9orf78 gene product has been provisionally designated C9orf78 pending further characterization.

Additional Information

Gene ID	51759
Other Names	Telomere length and silencing protein 1 homolog, Hepatocellular carcinoma-associated antigen 59, C9orf78, HCA59
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	C9orf78
Synonyms	HCA59 {ECO:0000303 PubMed:12097419}
Function	Plays a role in pre-mRNA splicing by promoting usage of the upstream 3'-splice site at alternative NAGNAG splice sites; these are sites featuring alternative acceptor motifs separated by only a few nucleotides (PubMed: 35241646). May also modulate exon inclusion events (PubMed: 35241646). Plays a role in spliceosomal remodeling by displacing WBP4 from SNRNP200 and may act to inhibit SNRNP200 helicase activity (PubMed: 35241646). Binds U5 snRNA (PubMed: 35241646). Required for proper chromosome segregation (PubMed: 35167828). Not required for splicing of shelterin components (PubMed: 35167828).
Cellular Location	Nucleus. Chromosome, centromere. Note=Dispersed throughout the nucleus during interphase (PubMed:35167828). Colocalizes with microtubule attachment sites at centromeres following mitotic checkpoint activation (PubMed:35167828).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.