

GPR177 Rabbit pAb

GPR177 Rabbit pAb Catalog # AP55975

Product Information

Application WB, IHC-P, IHC-F, IF

Primary Accession Q5T9L3

Reactivity Pig, Mouse, Dog, Horse

Host Rabbit
Clonality Polyclonal
Calculated MW 62253
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human GPR177

Epitope Specificity 351-450/541

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. **SUBCELLULAR LOCATION** Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasmic

vesicle membrane; Multi-pass membrane protein.

SIMILARITY Belongs to the wntless family.

SUBUNIT Interacts with WNT3A. Interacts with WNT1, WNT3 and WNT5A (By similarity). **Important Note** This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Additional Information

Gene ID 79971

Other Names Protein wntless homolog, Integral membrane protein GPR177, Protein

evenness interrupted homolog, EVI, Putative NF-kappa-B-activating protein

373, WLS, C1orf139, GPR177

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name WLS

Synonyms C1orf139, GPR177


Function Regulates Wnt proteins sorting and secretion in a feedback regulatory

mechanism. This reciprocal interaction plays a key role in the regulation of expression, subcellular location, binding and organelle-specific association of Wnt proteins (PubMed:34587386). Plays also an important role in establishment of the anterior-posterior body axis formation during development (By similarity).

Cellular Location

Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Early endosome membrane; Multi-pass membrane protein. Note=Co-localizes with the adaptin AP2A2 at distinct punctae.

Images




Sample: Brain (Mouse) Lysate at 40 ug

Primary: Anti-GPR177(AP55975) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000

dilution


Predicted band size: 62 kD Observed band size: 51 kD



Tissue/cell: rat kidney tissue;4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Blocking buffer (normal goat serum,C-0005) at 37 ∩ for 20 min;

Incubation: Anti-GPR177 Polyclonal Antibody, Unconjugated(AP55975) 1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated(AP55975-Cy3)used at 1:200 dilution for 40 minutes at 37°C. DAPI(5ug/ml,blue,C-0033) was used to stain the cell nuclei



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-GPR177 Polyclonal Antibody, Unconjugated(AP55975) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and

DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.