

HEV Capsid protein Rabbit pAb

HEV Capsid protein Rabbit pAb

Catalog # AP56006

Product Information

Application	WB, IHC-P, IHC-F, IF
Predicted	Hepatitis E Virus
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from Hepatitis E Virus Capsid protein
Epitope Specificity	301-400/660
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Virion (Probable). Host cytoplasm. Host cell surface. Note=Initially cotranslationally translocated into the ER from where it is retrotranslocated to the cytoplasm. A fraction is also observed on the cell surface.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The Hepatitis E virus is the causative agent of Hepatitis E. Its taxonomic name is Orthohepevirus A. The viral genome a single-strand of positive-sense RNA that is approximately 7200 bases in length. It encodes 3 proteins (O1, O2, O3), two of which are polyproteins, that is, they are cleaved into fragments which carry out the actual functions of the virus. The O1 protein consists of 7 such fragments, namely Met (Methyltransferase), Y (Y-domain), Plp (Papain like protease), V (proline-rich variable region), X (X-domain, macro-domain), Hel (Helicase), and RdRp (RNA dependent RNA polymerase). The Pvx domain is a fusion protein consisting of the Plp, V and X domains. The O3 protein is encoded by a single open reading frame (ORF3). The O2 protein encodes the capsid, which is composed of 3 domains, namely the shell domain (S) and two protruding domains (P1, P2). Numbers in the figure indicate positions in the RNA sequence.

Additional Information

Other Names	Capsid protein, Protein ORF2, pORF2, ORF2
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.