

FAM49B Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP56057

Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9NUQ9
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36748
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FAM49B
Epitope Specificity	251-324/324
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the FAM49 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Made up of nearly 146 million bases, chromosome 8 encodes about 800 genes. Translocation of portions of chromosome 8 with amplifications of the c-Myc gene are found in some leukemias and lymphomas, and typically associated with a poor prognosis. Portions of chromosome 8 have been linked to schizophrenia and bipolar disorder. Trisomy 8, also known as Warkany syndrome 2, most often results in early miscarriage but is occasionally seen in a mosaic form in surviving patients who suffer to a varying degree from a number of symptoms including retarded mental and motor development, and certain facial and developmental defects. WRN is a DNA helicase encoded by chromosome 8 and shown defective in those with the early aging disorder Werner syndrome. Chromosome 8 is also associated with Pfeiffer syndrome, congenital hypothyroidism and Waardenburg syndrome. The FAM49B gene product has been provisionally designated FAM49B pending further characterization.

Additional Information

Gene ID	51571
Other Names	CYFIP-related Rac1 interactor B, L1, CYRIB {ECO:0000303 PubMed:30250061, ECO:0000312 HGNC:HGNC:25216}
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Format	0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	CYRIB {ECO:0000303 PubMed:30250061, ECO:0000312 HGNC:HGNC:25216}
-------------	--

Function	Negatively regulates RAC1 signaling and RAC1-driven cytoskeletal remodeling (PubMed: 30250061 , PubMed: 31285585). Regulates chemotaxis, cell migration and epithelial polarization by controlling the polarity, plasticity, duration and extent of protrusions. Limits Rac1 mediated activation of the Scar/WAVE complex, focuses protrusion signals and regulates pseudopod complexity by inhibiting Scar/WAVE- induced actin polymerization (PubMed: 30250061). Protects against Salmonella bacterial infection. Attenuates processes such as macropinocytosis, phagocytosis and cell migration and restrict sopE- mediated bacterial entry (PubMed: 31285585). Also restricts infection mediated by Mycobacterium tuberculosis and Listeria monocytogenes (By similarity). Involved in the regulation of mitochondrial dynamics and oxidative stress (PubMed: 29059164).
-----------------	--

Cellular Location	Membrane; Lipid-anchor. Mitochondrion
--------------------------	---------------------------------------

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.