10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FBXO10 Rabbit pAb

FBXO10 Rabbit pAb
Catalog # AP56088

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9UK96
Reactivity	Mouse
Predicted	Rat, Human, Rabbit, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	105195
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FBXO10
Epitope Specificity	21-120/956
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Members of the F-box protein family, such as FBXO10, are characterized by an approximately 40-amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM, Mar 2008]

Additional Information

Gene ID	26267
Other Names	F-box only protein 10, FBXO10, FBX10, PRMT11
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	FBXO10
Synonyms	FBX10, PRMT11

Function	Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex. Mediates the ubiquitination and degradation of BCL2, an antiapoptotic protein, thereby playing a role in apoptosis by controlling the stability of BCL2. Targets also the receptor for advanced glycation end products RAGE for ubiquitination and subsequent lysosomal degradation (PubMed:28515150). Directly controls HGAL/GCSAM ubiquitination and degradation and thereby decreases BCR signaling (PubMed:31570756).
Cellular Location	Cytoplasm

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FBXO10) Polyclonal Antibody, Unconjugated (AP56088) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.