10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FITM1 Rabbit pAb

FITM1 Rabbit pAb
Catalog # AP56116

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	A5D6W6
Reactivity	Rat, Human
Predicted	Dog, Mouse, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32207
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FITM1
Epitope Specificity	1-100/292
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	FIT1 belongs to an evolutionarily conserved family of proteins involved in fat storage (Kadereit et al., 2008 [PubMed 18160536]).[supplied by OMIM, May 2008]

Additional Information

Gene ID	161247
Other Names	Fat storage-inducing transmembrane protein 1 {ECO:0000255|HAMAP-Rule:MF_03229, ECO:0000303|PubMed:18160536}, Fat-inducing protein 1 {ECO:0000255|HAMAP-Rule:MF_03229}, FITM1 {ECO:0000255|HAMAP-Rule:MF_03229, ECO:0000312|HGNC:HGNC:33714}
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	FITM1 {ECO:0000255|HAMAP-Rule:MF_03229, ECO:0000312|HGNC:HGNC:33714}

Function	Plays an important role in the formation of lipid droplets (LDs) which are storage organelles at the center of lipid and energy homeostasis (By similarity) (PubMed:18160536). Directly binds to diacylglycerol (DAGs) and triacylglycerol (By similarity).
Cellular Location	Endoplasmic reticulum membrane {ECO:0000255|HAMAP-Rule:MF_03229, ECO:0000269|PubMed:18160536}; Multi- pass membrane protein {ECO:0000255|HAMAP-Rule:MF_03229}
Tissue Location	Primarily expressed in heart and skeletal muscle.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Protein: 293T(human) lysate at 40ug;
Primary: rabbit Anti-FITM1 (AP56116) at 1:300;
Secondary: HRP conjugated Goat-Anti-rabbit IgG(AP56116-HRP) at 1: 5000;
Predicted band size: 32 kD
Observed band size: 28 kD



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 ∑ C for 30min; Antibody incubation with (Fat inducing transcript; FITM1) Polyclonal Antibody, Unconjugated (AP56116) at 1:400 overnight at 4 ∑ C, followed by a conjugated secondary antibody (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.