10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FRZB/FRP-3 Rabbit pAb

FRZB/FRP-3 Rabbit pAb
Catalog # AP56170

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q92765
Predicted	Rat, Pig, Dog, Human, Mouse, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36254
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FRZB/FRP-3
Epitope Specificity	1-100/251
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a secreted protein that is involved in the regulation of bone development. Defects in this gene are a cause of female-specific osteoarthritis (OA) susceptibility. [provided by RefSeq, Apr 2010]

Additional Information

Gene ID	2487
Other Names	Secreted frizzled-related protein 3, sFRP-3, Frezzled, Fritz, Frizzled-related protein 1, FrzB-1, FRZB, FIZ, FRE, FRP, FRZB1, SFRP3
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	FRZB
Synonyms	FIZ, FRE, FRP, FRZB1, SFRP3

Function	Soluble frizzled-related proteins (sFRPS) function as modulators of Wnt signaling through direct interaction with Wnts. They have a role in regulating cell growth and differentiation in specific cell types. SFRP3/FRZB appears to be involved in limb skeletogenesis. Antagonist of Wnt8 signaling. Regulates chondrocyte maturation and long bone development.
Cellular Location	Secreted.
Tissue Location	Expressed primarily in the cartilaginous cores of the long bone during embryonic and fetal development and in the appendicular skeleton (6-13 weeks). At 13 weeks of gestation, transcripts were present in early chondroblasts of the tarsal bones of the foot, the carpal bones of the hands and the epiphysis of long bones. Highly expressed in placenta and heart, followed by brain, skeletal muscle, kidney and pancreas. Weakly expressed in lung and liver

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.