

ZSCAN2/ZNF854 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP56248

Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q7Z7L9</u>

Reactivity Rat, Pig, Bovine

HostRabbitClonalityPolyclonalCalculated MW69547Physical StateLiquid

Immunogen KLH conjugated synthetic peptide derived from human ZSCAN2/ZNF854

Epitope Specificity 101-200/614

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nuclea

SIMILARITY Belongs to the krueppel C2H2-type zinc-finger protein family.Contains 14

C2H2-type zinc fingers. Contains 1 SCAN box domain.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions The protein encoded by this gene contains several copies of zinc finger motif,

which is commonly found in transcriptional regulatory proteins. Studies in mice show that this gene is expressed during embryonic development, and specifically in the testis in adult mice, suggesting that it may play a role in regulating genes in germ cells. Alternative splicing of this gene results in several transcript variants encoding different isoforms. [provided by RefSeq,

Jul 2008]

Additional Information

Gene ID 54993

Other Names Zinc finger and SCAN domain-containing protein 2, Zinc finger protein 29

homolog, Zfp-29, Zinc finger protein 854, ZSCAN2, ZFP29, ZNF854

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC=1:100-500,IF=1:100-50

0,ELISA=1:5000-10000

Format 0.01M TBS(pH7.4) with 1% BSA, 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name ZSCAN2


Synonyms ZFP29, ZNF854

Function May be involved in transcriptional regulation during the post-meiotic stages

of spermatogenesis.

Cellular Location Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00187}.

Images



Sample:

MCF-7(Human) Cell Lysate at 30 ug

Primary: Anti-ZNF854 (AP56248) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at

1/20000 dilution

Predicted band size: 70 kD Observed band size: 70 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.