

Hemoglobin subunit delta Rabbit pAb

Hemoglobin subunit delta Rabbit pAb

Catalog # AP56276

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P02042
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	16055
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Hemoglobin subunit delta
Epitope Specificity	2-80/147
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The delta (HBD) and beta (HBB) genes are normally expressed in the adult: two alpha chains plus two beta chains constitute HbA, which in normal adult life comprises about 97% of the total hemoglobin. Two alpha chains plus two delta chains constitute HbA-2, which with HbF comprises the remaining 3% of adult hemoglobin. Five beta-like globin genes are found within a 45 kb cluster on chromosome 11 in the following order: 5'-epsilon--Ggamma--Agamma--delta--beta-3'. Mutations in the delta-globin gene are associated with beta-thalassemia. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	3045
Other Names	Hemoglobin subunit delta, Delta-globin, Hemoglobin delta chain, HBD
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information


Name	HBD
-------------	-----

Function	Involved in oxygen transport from the lung to the various peripheral tissues.
Tissue Location	Red blood cells.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Images



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HBD) Polyclonal Antibody, Unconjugated (AP56276) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.