

HHV1 gE Rabbit pAb

HHV1 gE Rabbit pAb

Catalog # AP56289

Product Information

Application	IHC-P, IHC-F, IF
Predicted	Herpes simplex virus Strain F.
Host	Rabbit
Clonality	Polyclonal
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from HHV1 gE
Epitope Specificity	461-552/552
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Host cytoplasm. Host nucleus, host nucleolus. Virion. Note=At early times in infection, colocalizes with PCNA and replication proteins in cell nuclei, before accumulating in the cytoplasm by 8 to 12 hours post-infection. The effects on the host cell are mediated by de novo-synthesized ICP34.5, the virion-derived population being either non-functional or present in very low amounts.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Herpes simplex type 1 (HSV-1) belongs to a family that includes HSV-2, Epstein-Barr virus (EBV) and Varicella zoster (chicken pox) virus amongst others. HSV-1 and HSV-2 are extremely difficult to distinguish from each other. Members of this family have a characteristic virion structure. The double stranded DNA genome is contained within an icosahedral capsid embedded in a proteinaceous layer (tegument) and surrounded by a lipid envelope, derived from the nuclear membrane of the last host, which is decorated with virus-specific glycoproteins spikes. These viruses are capable of entering a latent phase where the host shows no visible sign of infection and levels of infectious agent become very low. During the latent phase the viral DNA is integrated into the genome of the host cell.

Additional Information

Other Names	Envelope glycoprotein E, gE, gE-1, gE
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.